



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

STEM-CELLBANKER®

Cryopreservation Medium

- Serum-Free
- Chemically Defined
- GMP Manufactured
- FDA Master File registered



Cat # 11924 (previously [11890])

Qty: 100ml

Stability: Guaranteed 3 years from manufacturing date (see label)


Cell-Freezing:

For optimum results, cells for cryopreservation should be in log phase of growth. Similar or standard freezing protocols may be substituted.

1. Examine and make sure the cell culture is free of contamination, in healthy situation and proper confluency, etc.
2. Perform a cell count to determine the viability of cells
3. Gently pellet the cells by centrifugation (3 - 5 minutes at 1,000~2,000rpm, 4°C). Remove the supernatant by using an aspirator.
4. Gently suspend the cells with STEM-CELLBANKER® cryopreservation medium (1 ml for 5×10^5 - 5×10^6 cells).
5. Dispense the cell suspension in 1ml aliquots to cryopreservation vials that have been labeled with the cell line name, cell concentration, passage date and other essential information.
6. Place the vials directly in a -80°C for storage. If necessary, transfer the frozen vials to a liquid nitrogen storage tank after the vials have been frozen for at least 24 hours.
7. Optimum protocol may change with the cell types.

IMPORTANT: Optimum protocol may change with the cell types.

Procedure for Use:



AMSBIO | www.amsbio.com | info@amsbio.com

UK & Rest of the World 
184 Park Drive, Milton Park
Abingdon OX14 4SE.
T: +44 (0) 1235 828 200
F: +44 (0) 1235 820 482

North America 
1035 Cambridge Street,
Cambridge, MA 02141.
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

Europe 
Berenkoog 41,
1822 BH Alkmaar,
Netherlands
T: +31 (0) 72 8080244
F: +31 (0) 72 8080142

Switzerland 
Via Lisano 3,
(CP.683)
CH-6900
T: +41 (0) 91 604 55 22
F: +41 (0) 91 605 17 85

Germany 
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880

Thawing:

1. Remove the frozen cell from storage and quickly thaw in a 37°C shaking water bath.
2. Immediately dilute and gently mix each 1ml of cells with 10ml of complete cell culture medium.
3. Gently pellet the cells by centrifugation (3-5 minutes at 1,000 - 2,000rpm, 4°C).
Remove the supernatant by aspirator.
4. Gently suspend the cells with appropriate volume of complete cell culture medium and plate in a culture flask.
5. Continue the further culture procedures according to standard protocols.

Guarantee of Quality:

1. Manufactured in compliance with JPN, EU, US, and PIC/S GMP guidelines
2. Bacterial contamination free - Product has been tested and confirmed to be free of bacteria, fungi and mycoplasma.
3. Chemical Analysis: pH (7.0 to 8.5 at room temperature) Endotoxin (<0.25 EU/mL)
4. Performance test - Cell viability above 80% (JM404, SK-007) is guaranteed.

Storage of STEM-CELLBANKER®:

1. STEM-CELLBANKER® should be stored at 2-8°C or below -20°C
2. For long-term storage STEM-CELLBANKER® can be frozen. Repeated freezing and thawing may impair the quality of the product; it is recommended that STEM-CELLBANKER® is aliquoted before freezing.

Disclaimer:

STEM-CELLBANKER® GMP grade is not by itself a pharmaceutical. Therefore, no warranty, express or implied, is made as to the fitness and suitability of this product for any particular purpose and/or merchantability unless the use is intended for research.

Product Range:

Code	Description	Pack Size
11884	CELLBANKER® 1 - Serum Containing	20 ml
11911	CELLBANKER® 1 - Serum Containing	4 x 20 ml
11910	CELLBANKER® 1 - Serum Containing	100 ml
11914	CELLBANKER® 2 - Serum Free	100 ml
11922	STEM-CELLBANKER® - GMP	20 ml
11894	STEM-CELLBANKER® - GMP	4 x 20 ml
11924	STEM-CELLBANKER® - GMP	100 ml
13925	STEM-CELLBANKER® - GMP - DMSO Free	20 ml
11894F	STEM-CELLBANKER® - GMP - DMSO Free	4 x 20 ml
13926	STEM-CELLBANKER® - GMP - DMSO Free	100 ml
11936	STEM-CELLBANKER® EX - GMP	100 ml
11918	CELLOTION cell wash solution	100 ml

AMSBIO | www.amsbio.com | info@amsbio.com

UK & Rest of the World 
184 Park Drive, Milton Park
Abingdon OX14 4SE.
T: +44 (0) 1235 828 200
F: +44 (0) 1235 820 482

North America 
1035 Cambridge Street,
Cambridge, MA 02141.
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

Europe 
Berenkoog 41,
1822 BH Alkmaar,
Netherlands
T: +31 (0) 728080244
F: +31 (0) 728080142

Switzerland 
Via Lisano 3,
(CP.683)
CH-6900
T: +41 (0) 91 604 55 22
F: +41 (0) 91 605 17 85

Germany 
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880

Citations:

Leung, S. T., Overschmidt, B., Allickson, J., Atala, A., Taghizadeh, R. R., Cetrulo, K., ... & Murphy, S. V. (2018). Review of Processing Technology and Techniques for Perinatal Stem Cells Banking and Clinical Applications. In Perinatal Stem Cells (pp. 337-355).
Miki, T., Vazquez, L., Yanuaria, L., Lopez, O., Garcia, I. M., Ohashi, K., & Rodriguez, N. S. (2019). Induced Pluripotent Stem Cell Derivation and Ex Vivo Gene Correction Using a Mucopolysaccharidosis Type 1 Disease Mouse Model. Stem Cells International, 2019.
Osaki, T., Uzel, S.G.M. & Kamm, R.D. On-chip 3D neuromuscular model for drug screening and precision medicine in neuromuscular disease. Nat Protoc (2020). https://doi.org/10.1038/s41596-019-0248-1
Skorik, C., Mullin, N. K., Shi, M., Zhang, Y., Hunter, P., Tang, Y., Hilton, B., & Schlaeger, T. M. (2020). Xeno- free reprogramming of peripheral blood mononuclear erythroblasts on laminin-521. Current Protocols in Stem Cell Biology, 52, e103. doi: 10.1002/cpsc.103
Ballantyne, M., Woodcock, M., Doddamani, D., Hu, T., Taylor, L., Hawken, R. J., & McGrew, M. J. (2021). Direct allele introgression into pure chicken breeds using Sire Dam Surrogate (SDS) mating. Nature communications, 12(1), 1-10.
Innes, J. A., Lowe, A. S., Fonseca, R., Aley, N., El-Hassan, T., Constantinou, M., ... & Brandner, S. (2021). Phenotyping clonal populations of glioma stem cell reveals a high degree of plasticity in response to changes of microenvironment. Laboratory Investigation, 1-13.
Zhang, K., Erkan, E. P., Jamalzadeh, S., Dai, J., Andersson, N., Kaipio, K., ... & Vähärautio, A. (2022). Longitudinal single-cell RNA-seq analysis reveals stress-promoted chemoresistance in metastatic ovarian cancer. Science advances, 8(8), eabm1831.

Cells tested:

Cell type	Description	Post Thaw Cell Viability
201B7	Human iPS cell	90
129SV	Mouse ES cell	90
P3/x63-Ag8.U1	Murine myeloma cell	90
2D-8	Murine hybridoma	90
YAC-1	Murine lymphoblast	90
NBM-Lu	Normal newborn murine fibroblast cell line	90
Feline PBMC	Feline peripheral blood mononuclear cells	80
Canine PBMC	Canine peripheral blood mononuclear cells	90
Jurkat	Human T-cell line	80
SK007	Human B-cell line	90
K562	Human Caucasian chronic myelogenous leukaemia cells	90
HeLa	Human uterine cervical carcinoma cell	90
HepG2	Human hepatocellular carcinoma cells	90
Caco-2	Human colonic adenocarcinoma cells	90
UE6E7-16	Human Mesenchymal cells	90
UE7T-13	Human Mesenchymal stem cells	90

FOR RESEARCH USE ONLY

AMSBIO | www.amsbio.com | info@amsbio.com

UK & Rest of the World 
184 Park Drive, Milton Park
Abingdon OX14 4SE.
T: +44 (0) 1235 828 200
F: +44 (0) 1235 820 482

North America 
1035 Cambridge Street,
Cambridge, MA 02141.
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

Europe 
Berenkoog 41,
1822 BH Alkmaar,
Netherlands
T: +31 (0)72 8080244
F: +31 (0)72 8080142

Switzerland 
Via Lisano 3,
(CP.683)
CH-6900
T: +41 (0) 91 604 55 22
F: +41 (0) 91 605 17 85

Germany 
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880