



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Ready-to-Use Dog Mixed Tissue Western Blot (custom made)

Catalog #: DW-CUS-2

Quantity: 1 blot

Storage Conditions: Store in a sealed bag at 4°C. Lot is good for several months.

Applications: Dog mixed tissue Western blot can be used to:

- Analyze protein expression patterns in 15 different Dog tissues simultaneously.
- Determine size and relative abundance of specific protein in different tissues

Quality Control: Every step of preparation of the blot, from harvesting tissues and extraction of protein to the blotting, is carefully monitored to ensure the superior quality and performance.

- Blot is made from high quality tissue lysates that is treated with a cocktail of mammalian protease inhibitors. The quality of protein as indicated by the absence of smear (no degradation) and sharpness and resolution of protein bands is verified by denatured SDS-PAGE with Coomassie blue staining.
- The efficiency of transfer is routinely checked by staining proteins on the membrane with reversible Ponceau staining.

Description: The total protein (tissue lysate; 50µg each) was extracted from freshly dissected dog tissues in lysis buffer containing a cocktail of 6 mammalian protease inhibitors, fractionated through standard (20x20cm) 12% SDS-denaturing polyacrylamide gel electrophoresis (SDS-PAGE), and blotted onto PVDF membrane.

To determine the size of the specific protein, a pre-stained molecular size protein marker is included in each blot. The protein bands are 250 KD, 150KD, 100 KD, 75 KD, 50 KD, 37 KD, 25 KD, 20 KD, and 15 KD.

Tissues on the blot: tissue lysates of 15 dog tissues selected by the end user

Prior to Use: Immerse blot in 100% methanol for few seconds then in water for 2 minutes before blocking the membrane. If blot is accidentally dry and showing white spots, immerse in methanol then in water.

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41(0) 91 604 55 22
F: +41(0) 91 605 17 85