



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

User's Manual and Instructions

IsHyb In Situ Hybridization (ISH) Kit

Catalog Number: K2191020, K2191050

Introduction

In situ hybridization (ISH) is a powerful method for gene expression analysis. Biochain is proud to launch a high-quality, fully customizable in situ hybridization kit for gene related studies. Biochain's IsHyb In Situ Hybridization Kit provides the reagents, including hybridization buffer, NBT/BCIP color development solution and washing buffer solutions, for pre-hybridization, hybridization, and post hybridization treatments.

Features

- microRNA detection
- Non-isotopic, colorimetric detection of gene expression
- RNase-free, ISH-optimized reagents
- Can be used with either oligonucleotide or ribonucleotide probes
- Tested by Digoxingenin labeled oligonucleotide probes
- Optimization of ISH assay conditions
- ISH detection in tissue section and/or tissue microarray slides
- Full technical support

Applications

- Diagnostic determination and clinical decision-making
- Gene mutation studies
- Protein expression and function analysis

Description

Biochain's ISH kit provides researchers a method doing in situ hybridization with nonradioactive probes. Compared with radioactive methods, the sensitivity of this kit is the same while the procedure is simplified. This kit is suitable for digoxingenin labeled probes combined with an alkaline phosphatase color development system. Using this kit, all disadvantages of working with radioactive materials can be avoided.

Quality control


In order to ensure the quality of IsHyb ISH kit, a pre-test procedure is performed for each lot of the kits. The positive result is detected with a digoxingenin labeled Has-mirRNA 122a cDNA probe on a previous positive frozen tissue panel slide.

Contents

Item	K2191020 Amount	K2191020 Part No.	K2191050 Amount	K2191050 Part No.
Prehybridization solution	2.0 ml	K2191020 -1	5 ml	K2191050 -1
Hybridization solution	2.0 ml	K2191020 -2	5 ml	K2191050 -2
NBT	120 µl	K2191020 -3	300 µl	K2191050 -3
BCIP	60 µl	K2191020 -4	150 µl	K2191050 -4
20 x SSC buffer	20.0 ml	K2191020 -5	50 ml	K2191050 -5
10x Alkaline phosphatase buffer	20.0 ml	K2191020 -6	50 ml	K2191050 -6
Anti-Dig Antibody	10 µl	K2191020 -7	25 µl	K2191050 -7
100 x Blocking solution	0.1 ml	K2191020 -8	0.25 ml	K2191050 -8

K2191020 is designed for hybridization of 20 slides, K2191050 is for 50 slides.

Buffer required but not included in the kit 1x PBS solution, pH 7.4: 81 ml 0.2 M Na₂HPO₄ 19 ml 0.2 M NaH₂PO₄ 18 g NaCl Add distilled water to 2,000 ml.



UK & Rest of World
184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America
23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio
info@amsbio.com
www.amsbio.com
AMS Biotechnology

Protocol

Preparation of frozen sections or paraffin sections

A: Pre-Treatment of Sections

- 1 Warm frozen slides to room temperature and dry at 50°C for 15 minutes
- 2 Paraffin slides: deparaffinization and rehydration through the following procedure:
 - 1) Deparaffinization: Soak tissue slides with xylene twice, each time for 15 minutes
 - 2) Rehydration: Let slides go through graded ethanol: 100%I, 100%II, 95%, 90%, 80%, 70%, and distilled water in the end. 5 minutes for each concentration.

Note: Baking the paraffin sections at 60°C for 1-2 hours is recommended if tissues drop off during following procedure

B: Hybridization

- 1 Fix the slides with 4% paraformaldehyde in DEPC-PBS at room temperature for 20 minutes.
- 2 Wash the slides twice with DEPC-PBS at room temperature for 5 minutes.
- 3 Treat the slides with 10 µg/ml Proteinase K at 37°C for 8-15 minutes.
- 4 Wash the slides once in DEPC-PBS at room temperature for five minutes. Fix them again with 4% paraformaldehyde in DEPC-PBS for 15 minutes. Rinse the slides once with DEPC-water.
- 5 Pre-hybridize the slides with ready-to-use **pre-hybridization solution** for 3-4 hours at **50°C**.
- 6 Hybridize the slides ready-to-use **hybridization solution** + Probe (Digoxigenin labeled probe: 2-4 ng/ul in hybridization solution) for 12-16 hours at 45°C.

Note: i) The volume of the pre-hyb, hyb., and blocking solutions should be just the amount that can cover the tissue section part on the slide. ii) Use an ImmEdge Pen (purchased from Vector) circle the tissue, then a boundary will be formed, then several drops of solution will be enough to cover the tissue. iii) During pre-hyb and hyb, in order to avoid the evaporation, put water soaked paper towel in a plastic box, and put any kind of rack on the top of paper towel, and put the slides on the rack, cover the plastic box then put the box in an incubator.


C: Washing Steps

- 1 Wash the slides with 2 x SSC for 10 minutes at 45°C.
- 2 Wash the slides with 1.5xSSC for 10 minutes at 45°C.
- 3 Wash the slides with 0.2xSSC twice at 37°C, 20 minutes for each time.
- 4 Incubate slides with 1 x **blocking solution** (dilute 100x blocking solution to 1x with 1x PBS) for 1 hour at room temperature. For some probes, the longer incubation will give the lower background.

D: Antibody Visualization of Digoxigenin

- 1 According to the sensitivity of the probe, incubate the slides with 1:100-1:1000 PBS diluted **AP-conjugated anti-digoxigenin antibody** for 1-4 hours or overnight at 4°C
- 2 Wash the slides with 1xPBS for 3 times at room temperature, 10 minutes each time.
- 3 Wash the slides with **1xAlkaline Phosphatase buffer** twice at room temperature, 5 minutes for each time.
- 4 According to the abundance of the target RNA, incubate the slides with solution, which is made by adding 6.6 µl NBT and 3.3 µl BCIP into 1 ml 1xAlkaline Phosphatase Buffer, in the dark for 2 to 20 hours at room temperature.
- 5 Discard the NBT/BCIP solution. Rinse the slides with distilled water
- 6 Counterstain if required.

Mount the slides.



UK & Rest of World

184 Milton Park, Abingdon
OX14 4SE, Oxon, UK
Tel: +44 (0) 1235 828 200
Fax: +44 (0) 1235 820 482

Switzerland

Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
Tel: +41 (0) 91 604 55 22
Fax: +41 (0) 91 605 17 85

Deutschland

Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
Tel: +49 (0) 69 779099
Fax: +49 (0) 69 13376880

North America

23591 El Toro Rd, Suite #180
Lake Forest, CA 92630
Tel: +1 800 987 0985
Fax: +1 949 265 7703

amsbio

info@amsbio.com

www.amsbio.com
AMS Biotechnology