



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse BLC Spleen cDNA

Code: MD-701-BLC

Quantity: 30 reactions

Storage Conditions: Store at -20°C. Suitable for one year from the date of purchase, if stored correctly.

Applications: The cDNA is primed with oligo dT primer and is ideal for gene expression analysis by PCR, characterization of alternative splicing of mRNA, gene cloning and target sequencing.

Quality Control: The PCR-Ready cDNA is functionally tested with the control primers in the recommended PCR reaction.

Description: The PCR-ready first strand cDNA is synthesized from high quality RNA isolated from tissue. This total RNA is isolated by modified guanidine thiocyanate method and treated with RNase-free DNase. The reverse transcribed is performed combining an oligo dT primer and reverse transcriptase, enabling the synthesis of full length cDNA up to 8.9kb. Use 1µl of cDNA for each PCR reaction.

The amplification conditions used for amplification of beta-actin as a positive control in a volume of 100µl:

Components	Volume (µl /reaction)	Final concentration
cDNA	1	
Polymerase reaction buffer	10	10X
MgCl ₂ , 25mM	7.8	2nM
Nucleotide Mix, 10mM	2.0	0.2mM
Upstream beta-Actin primer	1	100µM
Downstream beta-Actin primer	1	100µM
Taq Polymerase (5 units)	1	
Water	76.2	-
Total volume	100 µl	

Program of amplification:

	Denaturation	25 CYCLES			Final Extension	Hold
		Denaturation	Annealing	Extension		
Time	2 min.	1 min.	1 min.	2 min.	5 min.	-
Temperature	94 °C	94 °C	60 °C	72 °C	72 °C	4°C

Intended use of the products: All products sold by AMSBIO are intended for research use only and must not be used for medical diagnostic or drug purposes. AMSBIO products may not resell, modified for resale or used to manufacture commercial products without written approval from AMSBIO.

Limited product warranty: AMSBIO has a total quality assurance policy. If any product does not meet the specification stated in the product data sheet will be replaced at no charge. This warranty limits AMSBIO liability to the replacement of the product only. No other warranties of any kind, express or implied, including without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by AMSBIO. AMSBIO shall not be liable for any incidental, in direct, special or consequential damages, even if it is aware of the possibility of such damages.

AMSBIO | www.amsbio.com | info@amsbio.com



UK & Rest of the World
184 Park Drive, Milton Park
Abingdon OX14 4SE, UK
T: +44 (0)1235 828 200
F: +44 (0) 1235 820 482



North America
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218



Germany
Bockenheimer Landstr. 17/19
60325 Frankfurt/Main
T: +49 (0) 69 779099
F: +49 (0) 69 13376880



Switzerland
Centro Nord-Sud 2E
CH-6934 Bioggio-Lugano
T: +41 (0) 91 604 55 22
F: +41 (0) 91 605 17 85