

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Data Sheet

#### Ready-to-Use Mini-Pig Multiple Tissue Western Blot

Catalog #: NW-MT-1 Quantity: 1 blot

**Storage Conditions:** Store in a sealed bag at 4oC. It is good for several months.

Applications: Pig multiple tissues Western blot can be used to:

- Analyze protein expression pattern in 15 different rabbit tissues simultaneously.
- Determine size and relative abundance of specific protein in different tissues

**Quality Control:** Every step of preparation of the blot, from harvesting tissues and extraction of protein to the blotting, is carefully monitored to ensure the superior quality and performance.

- Blot is made from high quality tissue lysates that is treated with a cocktail of mammalian protease inhibitors. The quality of protein as indicated by the absence of smear (no degradation) and sharpness and resolution of protein bands is verified by denatured SDS-PAGE with Coomassie blue staining.
- The efficiency of transfer is routinely checked by staining proteins on the membrane with reversible Ponceau staining.
- The integrity of the blotted protein is tested by immunoblotting of  $\beta$ -actin.

**Description:** The total protein (tissue lysate; 75µg each) was extracted from freshly dissected Gottingen minipig tissues in lysis buffer containing a cocktail of 6 mammalian protease inhibitors, fractionated through standard (20x20cm) 12% SDS-denaturing polyacrylamide gel electrophoresis (SDS-PAGE), and blotted onto PVDF membrane.

To determine the size of the specific protein, a pre-stained molecular size protein marker is included in each blot. The protein bands are 250 KD, 150KD, 100 KD, 75 KD, 50 KD, 37 KD, 25 KD, 20 KD, 15 KD, and 10KD.

**Tissues on the blot:** Tissue lysates were loaded in the following order from the marker: (1) Brain, (2) stomach, (3) intestine, (4) colon, (5) liver, (6) lung (7) kidney, (8) Heart, (9) ovary, (10) skeletal muscles, (11) spleen, (12) testis, (13) thymus, (14) skin, and (15) pancreas.

**Prior to Use:** Immerse blot in 100% methanol for few seconds then in water for 2 minutes before blocking the membrane. If blot is accidentally dry and showing white spots, immerse in methanol then in water.

Intended use of the products: All products sold by AMSBIO are intended for research use only and must not be used for medical diagnostic or drug purposes. AMSBIO products may not resell, modified for resale or used to manufacture commercial products without written approval from AMSBIO. Limited product warranty: AMSBIO has a total quality assurance policy. If any product does not meet the specification stated in the product data sheet will be replaced at no charge. This warranty limits AMSBIO liability to the replacement of the product only. No other warranties of any kind, express or implied, including without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by AMSBIO.

AMSBIO shall not liable for any incidental, in direct, special or consequential damages, even if it is aware of the possibility of such damages.

**Deutschland** 

**United States**