



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



**Ready-to-Use Rat WS Lymph nodes Paraffin Sections****Catalog #:** RP-703-WS**Quantity:** 10 slides**Storage Conditions:** Store slides at room temperature in a dry place or at 4°C in a sealed bag containing desiccant.

Sections are good for 1 year of the date of purchase if stored properly.

**Applications:** Rat paraffin tissue sections are ideal for:

- Cellular localization of the protein of interest in rat tissues by immunohistochemistry.
- Cellular localization of the mRNA of interest in rat tissues by *in situ* hybridization or *in situ* PCR using both radioactive and non-radioactive-labelled probes.

**Description:** Tissues are freshly harvested from Sprague Dawley rats, fixed in 10% buffered neutral formalin, decalcified, and processed for embedding in low temperature melting paraffin. Tissue blocks are cut at a thickness of 5-7 and mounted on slides treated for adherence. Each slide contains 1 to 4 sections depending on the size of the tissue in the block. All tissue preparation steps from harvesting to embedding in paraffin are done in RNase-, DNase-, and protease-free conditions.**Shipping:** Slides are shipped at room temperature in a slide mailer or slide box.**Prior to use:** Remove paraffin by dipping slides in xylene or any substitutive agent (3 changes 5 minute each), rehydrate in serial ethanol dilutions, and finally wash with PBS. For *in situ* hybridization use, make alcohol solutions and PBS in RNase-free water.