

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





GSTM3 Antibody - middle region : FITC (ARP53562_P050-FITC)

Data Sheet

| Duo divot Niverbon | ADDESS (2) DOSO FITC |
|-------------------------------|--|
| Product Number | ARP53562_P050-FITC |
| Product Page | www.avivasysbio.com/gstm3-antibody-middle-region-fitc-arp53562-p050-fitc.html |
| Name | GSTM3 Antibody - middle region : FITC (ARP53562_P050-FITC) |
| Protein Size (# AA) | 225 amino acids |
| Molecular Weight | 26kDa |
| Conjugation | FITC: Fluorescein Isothiocyanate |
| NCBI Gene Id | 2947 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Concentration | 0.5 mg/ml |
| Gene Full Name | Glutathione S-transferase mu 3 (brain) |
| Alias Symbols | GST5, GSTB, GTM3, GSTM3-3 |
| Peptide Sequence | Synthetic peptide located within the following region: SMFLGKFSWFAGEKLTFVDFLTYDILDQNRIFDPKCLDEFPNLKAFMCRF |
| Product Format | Liquid. Purified antibody supplied in 1x PBS buffer. |
| Reference | Ikeda,S., (2008) Am J. Gastroenterol. 103 (6), 1476-1487 |
| Description of Target | Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. GSTM3 is a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Mutations of this class mu gene have been linked with a slight increase in a number of cancers, likely due to exposure with environmental toxins. Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Mutations of this class mu gene have been linked with a slight increase in a number of cancers, likely due to exposure with environmental toxins. Publication Note: This RefSeq r |
| Protein Interactions | GSTM5; GSTM4; GSTM3; GSTM1; UBC; ESR1; BAG3; SUMO2; SH3GL2; UCHL5; GSTM2; MPG; |
| Reconstitution and Storage | All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding. |
| Datasheets/Manuals | Printable datasheet for anti-GSTM3 (ARP53562_P050-FITC) antibody |
| Blocking Peptide | For anti-GSTM3 (ARP53562_P050-FITC) antibody is Catalog # AAP53562 (Previous Catalog # AAPS32802) |
| Immunogen | The immunogen is a synthetic peptide directed towards the middle region of human GSTM3 |
| Uniprot ID | P21266 |
| Protein Name | Glutathione S-transferase Mu 3 |
| | |

| Protein Accession # | <u>NP_000840</u> |
|--|--|
| Purification | Affinity Purified |
| Nucleotide Accession# | <u>NM_000849</u> |
| Gene Symbol | GSTM3 |
| Predicted Species Reactivity | Human, Mouse, Rat, Cow, Dog, Goat, Guinea Pig, Horse, Pig, Rabbit |
| Application | WB |
| Predicted Homology Based on Immunogen Sequence | Cow: 79%; Dog. 79%; Goat: 75%; Guinea Pig. 79%; Horse: 77%; Human: 100%; Mouse: 86%; Pig. 85%; Rabbit: 90%; Rat: 85% |
| Image 1 | |

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use. Optimal conditions of its use should be determined by end users.

AVIVA SYSTEMS BIOLOGY

6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | info@avivasysbio.com