



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Bovine IgG Heavy and Light Chain Cross-Adsorbed Antibody

Goat Polyclonal

Conjugate FITC

Antigen Affinity Purified

Catalog No. A10-230F

Lot No. 2

APPLICATIONS	IHC, ICC, Flow Cyt, IF								
SPECIES REACTIVITY	Bovine. Minimum reactivity to human, mouse and rat								
AMOUNT	1 ml								
CONCENTRATION	0.5 mg/ml								
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt								
PHYSICAL STATE	Liquid								
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide								
FLUOROPHORE/PROTEIN	4.2								
ISOTYPE	IgG								
ORIGIN	USA								
PRODUCTION PROCEDURES	<p>Antiserum was cross adsorbed using human, mouse, and rat immunosorbents to remove cross reactive antibodies. The antibody to bovine IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to fluorescein isothiocyanate.</p> <p>Immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with bovine IgG and with light chains common to other bovine immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human, mouse, and rat IgG was detected. This antibody may cross react with IgG from other species.</p>								
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Immunohistochemistry</td><td>1:50 – 1:500</td></tr><tr><td>Immunocytochemistry</td><td>1:50 – 1:500</td></tr><tr><td>Flow Cytometry</td><td>1:50 – 1:200</td></tr><tr><td>Immunofluorescence</td><td>1:50 – 1:500</td></tr></table>	Immunohistochemistry	1:50 – 1:500	Immunocytochemistry	1:50 – 1:500	Flow Cytometry	1:50 – 1:200	Immunofluorescence	1:50 – 1:500
Immunohistochemistry	1:50 – 1:500								
Immunocytochemistry	1:50 – 1:500								
Flow Cytometry	1:50 – 1:200								
Immunofluorescence	1:50 – 1:500								
APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.								
ADDITIONAL INFO	<p>https://www.fortislife.com/p/A10-230F</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>								

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: May 14, 2024