



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rat IgG–Fc Fragment Cross–Adsorbed Antibody

F(ab')₂ Goat Polyclonal

Conjugate R–Phycoerythrin

Antigen Affinity Purified

Catalog No. A110–239PE

Lot No. 8

APPLICATIONS	Flow Cyt, IF				
SPECIES REACTIVITY	Rat. Minimum reactivity to human and mouse				
AMOUNT	1 ml				
CONCENTRATION	0.5 mg/ml				
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt				
PHYSICAL STATE	Liquid				
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide				
ISOTYPE	IgG				
ORIGIN	USA				
PRODUCTION PROCEDURES	<p>Antiserum was solid phase adsorbed to ensure class specificity. Antiserum was cross adsorbed using human and mouse immunosorbents to remove cross reactive antibodies. The antibody to rat IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')₂ fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules were removed. Fragments were conjugated to R–phycoerythrin (RPE).</p> <p>Immunoglobulin concentration was determined using Beer's Law where 1 mg/mL IgG has an A₂₈₀ of 1.4.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non–immunoglobulin serum proteins. Less than 1% cross reactivity to human and mouse IgG was detected. This antibody may cross react with IgG from other species.</p>				
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Flow Cytometry</td><td>1:50 – 1:200</td></tr><tr><td>Immunofluorescence</td><td>1:50 – 1:200</td></tr></table>	Flow Cytometry	1:50 – 1:200	Immunofluorescence	1:50 – 1:200
Flow Cytometry	1:50 – 1:200				
Immunofluorescence	1:50 – 1:200				
APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.				
ADDITIONAL INFO	<p>https://www.fortislifesciences.com/p/A110–239PE</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>				

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: March 12, 2024