

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

| Rat IgG Heavy<br>Antibody  | and Light Chain Cross-Adsorbed  |
|----------------------------|---|
| Goat Polyclonal            | Conjugate DyLight <sup>®</sup> 594  |
| ,<br>Antigen Affinity Puri |   |
| Catalog No. A110           | 0-305D4   |
| Lot No. 17                 |   |
| APPLICATIONS               | IHC, ICC, Flow Cyt, IF  |
| SPECIES REACTIVITY         | Rat. Minimum reactivity to bovine, chicken, human, mouse, rabbit and sheep  |
| AMOUNT                     | 1 ml  |
| CONCENTRATION              | 0.5 mg/ml   |
| STORAGE/SHELF LIFE         | 2 – 8°C / 1 year from date of receipt   |
| PHYSICAL STATE             | Liquid  |
| BUFFER                     | Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide  |
| FLUOROPHORE/PROTE          | IN 6.4  |
| ISOTYPE                    | IgG   |
| ORIGIN                     | USA   |
| PRODUCTION<br>PROCEDURES   | Antiserum was cross adsorbed using bovine, chicken, human, mouse, rabbit and sheep immunosorbents to remove cross reactive antibodies. The antibody to rat IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 594.   |
|                            | Immunoglobulin concentration was determined using Beer's Law where $1 mg/mL$ IgG has an A280 of 1.4.  |
|                            | By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgG and with<br>light chains common to other rat immunoglobulins. No antibody was detected against non-<br>immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, chicken, human,<br>mouse, rabbit and sheep IgG was detected. This antibody may cross react with IgG from<br>other species. |
| APPLICATIONS               | Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.   |
|                            | Immunohistochemistry 1:50 – 1:500   |
|                            | Immunocytochemistry 1:50 – 1:500  |
|                            | Flow Cytometry 1:50 – 1:200<br>Immunofluorescence 1:50 – 1:500  |
| APPLICATION NOTES          | Not all listed applications have been specifically tested by our laboratory.  |
| AITEICATION NOTES          |   |
|                            | DyLight® 594 is excited at 593 (in PBS) and emits at 618 (in PBS).  |
|                            | DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.  |
| ADDITIONAL INFO            | https://www.fortislife.com/p/A110-305D4<br>Use the link above to view SDS, a current list of citations, and other product specific information.   |
|                            |   |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: March 7, 2024



