

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## Rabbit IgM Cross-Adsorbed Antibody





Antigen Affinity Purified

Catalog No. A120-210F

Lot No. 11

**APPLICATIONS** IHC, ICC, Flow Cyt, IF

SPECIES REACTIVITY Rabbit. Minimum reactivity to human, mouse and rat

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 5.4
ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antiserum was solid phase adsorbed to ensure class specificity. Antiserum was cross adsorbed using human, mouse and rat immunosorbents to remove cross reactive antibodies. The antibody to rabbit IgM was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to fluorescein isothiocyanate (FITC).

Immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an

A280 of 1.4.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgM. Cross

reactivity with IgA and IgG is negligible. No antibody was detected against non-

immunoglobulin serum proteins. Less than 1% cross reactivity to human, mouse and rat IgM

was detected. This antibody may cross react with IgM from other species.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry1:50 - 1:500Immunocytochemistry1:50 - 1:500Flow Cytometry1:50 - 1:200Immunofluorescence1:50 - 1:500

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO https://www.fortislife.com/p/A120-210F

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: April 1, 2024

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com Orders: orders@fortislife.com • Support: technical@fortislife.com