



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rabbit IgG Heavy and Light Chain Highly Cross-Adsorbed Antibody

Goat Polyclonal Conjugate FITC

Antigen Affinity Purified

Catalog No. A120-501F

Lot No. 7

APPLICATIONS	IHC, ICC, Flow Cyt, IF								
SPECIES REACTIVITY	Rabbit. Minimum reactivity to human and mouse								
AMOUNT	1 ml								
CONCENTRATION	1 mg/ml								
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt								
PHYSICAL STATE	Liquid								
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide								
FLUOROPHORE/PROTEIN	5.7								
ISOTYPE	IgG								
ORIGIN	USA								
PRODUCTION PROCEDURES	<p>Antiserum was cross adsorbed using human and mouse immunosorbents to remove cross reactive antibodies. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to fluorescein isothiocyanate (FITC).</p> <p>Immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4.</p> <p>This antibody reacts specifically with rabbit IgG in immunohistochemistry and immunocytochemistry fluorescence assays. Non-specific binding to human and mouse tissue is negligible.</p>								
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Immunohistochemistry</td><td>1:100 – 1:500. For use with Formalin-Fixed Parrafin-Embedded Tissues</td></tr><tr><td>Immunocytochemistry</td><td>1:100 – 1:500</td></tr><tr><td>Flow Cytometry</td><td>1:50 – 1:200</td></tr><tr><td>Immunofluorescence</td><td>1:100 – 1:500. For use with cells</td></tr></table>	Immunohistochemistry	1:100 – 1:500. For use with Formalin-Fixed Parrafin-Embedded Tissues	Immunocytochemistry	1:100 – 1:500	Flow Cytometry	1:50 – 1:200	Immunofluorescence	1:100 – 1:500. For use with cells
Immunohistochemistry	1:100 – 1:500. For use with Formalin-Fixed Parrafin-Embedded Tissues								
Immunocytochemistry	1:100 – 1:500								
Flow Cytometry	1:50 – 1:200								
Immunofluorescence	1:100 – 1:500. For use with cells								
APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.								
ADDITIONAL INFO	<p>https://www.fortislife.com/p/A120-501F</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>								

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: July 24, 2024