

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



KIAA0460 Antibody





Goat Polyclonal

Antigen Affinity Purified Protein ID CAH70314.1

Catalog No. A300-031A-T GeneID 23248

Lot No. A300-031A-T-2

APPLICATIONS WB

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to KIAA0460 immobilized on solid

PROCEDURES support.

The epitope recognized by A300-031A-T maps to a region between residue 700 and 750 of

human KIAA0460 using the numbering given in entry CAH70314.1 (GenelD 23248).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280

nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:5,000

Immunoprecipitation Not recommended. Use goat anti-KIAA0460 antibody A300-032A.


ADDITIONAL INFO https://www.bethyl.com/product/A300-031A-T

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022



Detection of human KIAA0460 by western blot. Samples: Whole cell lysate (50 μ g) from HeLa and HEK293T cells prepared using NETN lysis buffer. Antibody: Affinity purified goat anti-KIAA0460 antibody A300-031A (lot A300-031A-2) used for WB at 0.2 μ g/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.