



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SA1 Antibody

Goat Polyclonal

Antigen Affinity Purified	RefSeq ID	NP_005853.2
Catalog No. A300-157A-T	Uniprot ID	Q8WVM7
Lot No. 4	GeneID	10274


APPLICATIONS	WB, IP, ICC						
SPECIES REACTIVITY	Human						
AMOUNT	10 µl						
CONCENTRATION	1000 µg/ml						
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt						
PHYSICAL STATE	Liquid						
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide						
ISOTYPE	IgG						
ORIGIN	USA						
PRODUCTION PROCEDURES	<p>Antibody was affinity purified using an epitope specific to SA1 immobilized on solid support.</p> <p>The epitope recognized by A300-157A-T maps to a region between residues 1150 and 1200 of human Stromal Antigen 1 using the numbering given in entry NP_005853.2 (GeneID 10274).</p> <p>Immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4.</p>						
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:2000 – 1:10,000</td></tr><tr><td>Immunoprecipitation</td><td>6 µg/mg lysate</td></tr><tr><td>Immunocytochemistry</td><td>1:500 – 1:2000</td></tr></table>	Western Blot	1:2000 – 1:10,000	Immunoprecipitation	6 µg/mg lysate	Immunocytochemistry	1:500 – 1:2000
Western Blot	1:2000 – 1:10,000						
Immunoprecipitation	6 µg/mg lysate						
Immunocytochemistry	1:500 – 1:2000						
ADDITIONAL INFO	<p>https://www.fortislife.com/p/A300-157A-T</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>						

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: May 10, 2024



Detection of human SA1 by western blot. *Samples:* Whole cell lysate (25 µg) from HeLa, HEK293T, Jurkat, MCF-7, and GaMG cells prepared using NETN lysis buffer. *Antibody:* Affinity purified goat anti-SA1 antibody (A300-157A lot 4) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.

Detection of human SA1 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 15% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-SA1 antibody (A300-157A lot 4) used for IP at 6 µg per reaction. SA1 was also immunoprecipitated by a previous lot of this antibody (A300-157A lot 3) and a second antibody against a different epitope of SA1 (A302-578A). For blotting immunoprecipitated SA1, A300-157A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



Localization of human SA1. *Sample:* HeLa cells that were extracted for 5 min. at 4C in 0.5% Triton in CSK buffer. *Antibody:* Affinity purified goat anti-SA1 (A300-157A) used at 1 µg/ml.