

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PNK1 Antibody

Antigen Affinity Purified

Liquid

IgG

USA

support.

Western Blot

Immunoprecipitation

Immunohistochemistry

TrEMBL entry Q96T60 (GeneID 11284).

Rabbit Polyclonal

Catalog No.

APPLICATIONS

SPECIES REACTIVITY

CONCENTRATION

PHYSICAL STATE

STORAGE/SHELF LIFE

Lot No.

AMOUNT

BUFFER

ISOTYPE

ORIGIN

PRODUCTION

PROCEDURES

APPLICATIONS

body					
nal					
ty Purified		Protein ID	Q96T60		
A300-257A-T		GenelD	11284		
A300-	-257A-T-2				
	WB, IHC				
VITY	Human				
	10 µl				
ON	1000 µg/ml				
F LIFE	2 - 8°C / 1 year from date of receipt				
E	Liquid				

recommended for FFPE tissue sections. **ADDITIONAL INFO** https://www.bethyl.com/product/A300-257A-T Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the guality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: June 6, 2022

1:2.000 - 1:10.000

Not recommended

Tris-citrate/phosphate buffer. pH 7 to 8 containing 0.09% Sodium Azide

Antibody was affinity purified using an epitope specific to PNK1 immobilized on solid

experimentally by the investigator. Prepare working dilution immediately before use.


The epitope recognized by A300-257A-T maps to a region between residues 100 and 150 of human polynucleotide kinase 3-prime-phosphatase using the numbering given in


Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

1:500 – 1:2.000. Epitope retrieval with citrate buffer pH 6.0 is

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.

2





Detection of human PNK1 by western blot. *Samples:* Whole cell lysate (50 μ g) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-PNK1 antibody A300-257A (lot A300-257A-2) used for WB at 0.1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.

Detection of human PNK1 by immunohistochemistry. *Sample:* FFPE section of human ovarian carcinoma. *Antibody:* Affinity purified rabbit anti-PNK1 (Cat. No. A300-257A Lot2) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* DAB