

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



### **FUS Antibody**

Rabbit Polyclonal

SPECIES REACTIVITY



Antigen Affinity Purified RefSeq ID NP\_004951.1

Catalog No. A300-293A-T Uniprot ID P35637 Lot No. 6 GeneID 2521

Human, Mouse

**APPLICATIONS** WB, IP, IHC

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Boyine

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to FUS immobilized on solid support.

The epitope recognized by A300–293A–T maps to a region between residues 400 and 450 of human fusion (involved in t(12:16) in malignant liposarcoma) using the numbering given

in SwissProt entry P35637 (GeneID 2521).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000Immunoprecipitation  $2 - 10 \mu g/mg$  lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE tissue sections.

IHC HUMAN CONTROLS Breast Carcinoma, Colon Carcinoma, Ovarian Carcinoma, Prostate Carcinoma, Skin Squamous

Cell Carcinoma

IHC MOUSE CONTROLS Renal Cell Carcinoma, Squamous Cell Carcinoma, Teratoma

ADDITIONAL INFO https://www.fortislife.com/p/A300-293A-T


Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: April 14, 2023


FUS Antibody A300-293A-T







Detection of human FUS by western blot. Samples: Whole cell lysate (25  $\mu$ g) from HeLa, HEK293T, Jurkat, A-549, and U2OS cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-FUS antibody (A300–293A lot 6) used for WB at 0.1  $\mu$ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

Detection of mouse FUS by western blot. Samples: Whole cell lysate (10 μg) from NIH 3T3, CT26, TCMK-1, and BW5147.3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-FUS antibody (A300-293A-6 lot 6) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 75 seconds.






purified rabbit anti–FUS antibody (A300–293A lot 6) used for IP at 6 µg per reaction. FUS was also immunoprecipitated by a previous lot of this antibody (A300–293A lot 5) and a second antibody against a different epitope of FUS (A300–294A). For blotting immunoprecipitated FUS, A300–293A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 seconds.



Detection of human FUS by immunohistochemistry. Sample: FFPE section of human ovarian carcinoma. Antibody: Affinity purified rabbit anti-FUS (A300-293A lot 6) used at of 1:5,000 (0.2µg/ml). Detection: DAB

FUS Antibody A300-293A-T



Detection of mouse FUS by immunohistochemistry. *Sample:* FFPE section of mouse spleen. *Antibody:* Affinity purified rabbit anti-FUS (A300–293A lot 6) used at 1:5,000 (0.2µg/ml). *Detection:* DAB