

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Ubiquitin Antibody



Antigen Affinity Purified Protein ID P62988
Catalog No. A300–317A–T GeneID 7311

Lot No. A300-317A-T-1

APPLICATIONS WB, IHC

SPECIES REACTIVITY Human, Mouse, Bovine

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat, D. melanogaster,

X. laevis, Chicken, Sheep, Dog, Horse, Rabbit, Guinea pig_10141, Pig, Cat, Gorilla and

Chimpanzee

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to Ubiquitin immobilized on solid

PROCEDURES support.

The epitope recognized by A300-317A-T maps to a region between residues 1 and 50 of

human Ubiquitin using the numbering given in entry P62988.(GenelD 7311).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:10,000

Immunohistochemistry 1:500 - 1:2,000. Epitope retrieval with Tris-EDTA pH 9.0 is

recommended for FFPE tissue sections.


ADDITIONAL INFO https://www.bethyl.com/product/A300-317A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022



Detection of Ubiquitin by western blotting of Products from an Ubiquitination Assay and Purified Ubiquitin. Samples: A) Ubiquitination assay was performed as described in Zhang et al., 2004, Mol. Cell. Biol. 24(24):10941–10953. B) Purified ubiquitin applied in decreasing amounts from 5 to 0.1 μg. Antibody: Affinity purified rabbit anti-Ubiquitin antibody (Cat. No. A300–317A) used at 0.4 μg/ml for western blot. Detection: Alkaline phosphatase conjugated secondary antibody and development with NBT/BCIP for 10 minutes (A and B).



Detection of human and mouse Ubiquitin by immunohistochemistry. Sample: FFPE sections of human breast carcinoma (left) and mouse teratoma (right). Antibody: Affinity purified rabbit anti-Ubiquitin (Cat. No. A300-317A Lot1) used at a dilution of 1:1,000 (1µg/ml). Detection: DAB