

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



ZBTB7/FBI-1 Antibody





Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_056982.1

Catalog No. A300-548A-T GeneID 51341

Lot No. A300-548A-T-3

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to ZBTB7/FBI-1 immobilized on

PROCEDURES solid support.

The epitope recognized by A300-548A-T maps to a region between residues 200 and 250

of human ZBTB7 (Factor Binding IST protein 1) using the numbering given in entry

NP 056982.1 (GeneID 51341).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 5 µg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A300-548A-T


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022






Detection of human ZBTB7/FBI-1 by western blot.

Samples: Whole cell lysate (50 μg) from HeLa and HEK293T cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-ZBTB7/FBI-1 antibody A300-548A (lot A300-548A-3) used for WB at 0.1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.

Detection of mouse ZBTB7/FBI-1 by western blot. Samples: Whole cell lysate (50 μg) from TCMK-1 and NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-ZBTB7/FBI-1 antibody A300-548A (lot A300-548A-3) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Detection of human ZBTB7/FBI-1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-ZBTB7/FBI-1 antibody A300-548A (lot A300-548A-3) used for IP at 3 μg per reaction. ZBTB7/FBI-1 was also immunoprecipitated by a previous lot of this antibody (lot A300-548A-2) and rabbit anti-ZBTB7/FBI-1 antibody A300-549A. For blotting immunoprecipitated ZBTB7/FBI-1, A300-548A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.