



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NUP214 Antibody

Rabbit Polyclonal

Antigen Affinity Purified	RefSeq ID	NP_005076.3
Catalog No. A300-716A	Uniprot ID	P35658
Lot No. A300-716A-3	GeneID	8021

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human
AMOUNT	100 µl
CONCENTRATION	200 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to NUP214 immobilized on solid support.

The epitope recognized by A300-716A maps to a region between residue 1250 and 1300 of human Nucleoporin 214 using the numbering given in entry NP_005076.3 (GeneID 8021).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate


ADDITIONAL INFO <https://www.fortislife.com/p/A300-716A>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: September 14, 2022



Detection of human NUP214 by western blot. *Samples:* Whole cell lysate (25 µg) from HeLa, K-562, RKO, LNCaP, and HEK293T cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-NUP214 antibody (A300-716A lot 3) used for WB at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human NUP214 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 5% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-NUP214 antibody (A300-716A lot 3) used for IP at 6 µg per reaction. NUP214 was also immunoprecipitated by a previous lot of this antibody (A300-716A lot 2) and a second antibody against a different epitope of NUP214 (A300-717A). For blotting immunoprecipitated NUP214, A300-716A was used at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.