

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



USP36/DUB1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_079366.2

Catalog No. A300-940A GeneID 57602

Lot No. A300-940A-2

APPLICATIONS WB, IP, IHC
SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

DRIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to USP36/DUB1 immobilized on solid support.

The epitope recognized by A300-940A maps to a region between residue 1070 and the C-terminus (residue 1121) of human Ubiquitin Specific Peptidase 36 (Deubiquinating Enzyme 1)

using the numbering given in entry NP_079366.2 (GeneID 57602).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,,000 - 1:5,000Immunoprecipitation $2 - 10 \mu g/mg$ lysate

Immunohistochemistry 1:2,000 - 1:10,000. Epitope retrieval with Tris-EDTA pH 9.0 is

recommended for FFPE tissue sections.

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

IHC HUMAN CONTROLS Breast Carcinoma, Colon Carcinoma, Non-Small Cell Lung Cancer, Ovarian Carcinoma, Prostate

Carcinoma

ADDITIONAL INFO https://www.bethyl.com/product/A300-940A


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush. PhD | Chief Scientific Officer

Date: June 21, 2019








Detection of Human USP36/DUB1 by Western Blot. Samples: Whole cell lysate (50 μg) from HeLa, Hep-G2, and U2OS cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-USP36/DUB1 antibody A300-940A (lot A300-940A-2) used for WB at 0.4 μg/ml. Detection: Chemiluminescence with an exposure time of 75 seconds.

Detection of human USP36/DUB1 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-USP36/DUB1 antibody A300–940A (lot A300–940A–2) used for IP at 6 μg per reaction. USP36/DUB1 was also immunoprecipitated by rabbit anti-USP36/DUB1 antibody BL4146. For blotting immunoprecipitated USP36/DUB1, A300–940A was used at 1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human USP36/DUB1 by immunohistochemistry. *Sample:* FFPE section of human colon carcinoma. *Antibody:* Affinity purified rabbit anti-USP36/DUB1(Cat. No. A300-940A Lot 2) used at a dilution of 1:5,000 (0.2µg/ml). *Detection:* DAB