



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UTP18/WDR50 Antibody

Rabbit Polyclonal

Antigen Affinity Purified	Protein ID	NP_057085.2
Catalog No. A301-552A-T	GeneID	51096
Lot No. A301-552A-T-2		

APPLICATIONS	IP
SPECIES REACTIVITY	Human
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to UTP18/WDR50 immobilized on solid support.

The epitope recognized by A301-552A-T maps to a region between residue 325 and 375 of human small subunit (SSU) processome component UTP18 (WD repeat domain 50) using the numbering given in entry NP_057085.2 (GeneID 51096).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot	Not recommended. Use rabbit anti-UTP18/WDR50 antibody A301-551A.
--------------	--

Immunoprecipitation	2 – 5 µg/mg lysate
---------------------	--------------------

ADDITIONAL INFO <https://www.bethyl.com/product/A301-552A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.
IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human UTP18/WDR50 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-UTP18/WDR50 antibody A301-552A (lot A301-552A-2) used for IP at 3 µg per reaction. UTP18/WDR50 was also immunoprecipitated by rabbit anti-UTP18/WDR50 antibody A301-551A. For blotting immunoprecipitated UTP18/WDR50, A301-551A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.