

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



STUB1/CHIP Antibody





Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_005852.2

Catalog No. A301-572A-T GeneID 10273

Lot No. A301-572A-T-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to STUB1/CHIP immobilized on solid

PROCEDURES support.

The epitope recognized by A301-572A-T maps to a region between residue 1 and 50 of human STIP1 homology and U-box containing protein 1 (carboxy terminus of Hsp70-interacting protein) using the numbering given in entry NP_005852.2 (GenelD 10273).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2.000 - 1:10.000

Immunoprecipitation 2 – 5 μg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A301-572A-T


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022





Detection of human and mouse STUB1/CHIP by western blot. Samples: Whole cell lysate from HeLa (5, 15 and 50 μ g), HEK293T (T; 50 μ g) and mouse NIH 3T3 (M; 50 μ g) cells. Antibodies: Affinity purified rabbit anti-STUB1/CHIP antibody A301-572A (lot A301-572A-1) used for WB at 0.04 μ g/ml. Detection: Chemiluminescence with exposure time of 30 seconds.

Detection of human STUB1/CHIP by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti–STUB1/CHIP antibody A301–572A (lot A301–572A–1) used for IP at 3 μ g/mg lysate. STUB1/CHIP was also immunoprecipitated by rabbit anti–STUB1/CHIP antibody A301–573A, which recognizes a downstream epitope. For blotting immunoprecipitated STUB1/CHIP, A301–572A was used at 0.1 μ g/ml. Detection: Chemiluminescence with exposure time of 30 seconds.