



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Myo18A/Myosin XVIII A Antibody

Rabbit Polyclonal

Antigen Affinity Purified	Protein ID	NP_510880.2
Catalog No. A301-598A-T	GeneID	399687
Lot No. A301-598A-T-1		

APPLICATIONS	IP
SPECIES REACTIVITY	Human
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to Myo18A/Myosin XVIII A immobilized on solid support.

The epitope recognized by A301-598A-T maps to a region between residue 2004 and 2054 of human myosin 18A using the numbering given in entry NP_510880.2 (GeneID 399687).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot	Not recommended. A301-598A has not performed satisfactorily when used for WB of Myo18A/Myosin XVIII A in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of Myo18A/Myosin XVIII A.
--------------	--


Immunoprecipitation	2 – 5 µg/mg lysate
---------------------	--------------------

ADDITIONAL INFO <https://www.bethyl.com/product/A301-598A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human Myo18A/Myosin XVIII A by western blot of immunoprecipitates. *Samples:* RIPA whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-Myo18A/Myosin XVIII A antibody A301-598A used for IP at 3 µg/mg lysate. Myo18A/Myosin XVIII A was also immunoprecipitated by rabbit anti-Myo18A/Myosin XVIII A antibody A301-597A, which recognizes an upstream epitope. For blotting immunoprecipitated Myo18A/Myosin XVIII A, A301-597A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 1 second.