



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

iASPP Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_006654.2

Catalog No. A301-737A-T

GenelD 10848

Lot No. A301-737A-T-2

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
AMOUNT	10 µl
CONCENTRATION	200 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to iASPP immobilized on solid support.

The epitope recognized by A301-737A-T maps to a region between residue 75 and 125 of human inhibitor of apoptosis stimulating protein of p53 using the numbering given in entry NP_006654.2 (GenelD 10848).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

ADDITIONAL INFO <https://www.bethyl.com/product/A301-737A-T>


Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human and mouse iASPP by western blot.

Samples: Whole cell lysate from HEK293T (50, 15, 5 µg), HeLa (50, 15 µg), and mouse NIH 3T3 (50 µg) cells prepared using NETN lysis buffer. **Antibody:** Affinity purified rabbit anti-iASPP antibody A301-737A (lot A301-737A-2) used for WB at 0.1 µg/ml. **Detection:** Chemiluminescence with an exposure time of 30 seconds.



Detection of human iASPP by western blot of immunoprecipitates.

Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. **Antibodies:** Affinity purified rabbit anti-iASPP antibody A301-737A (lot A301-737A-2) used for IP at 6 µg per reaction. iASPP was also immunoprecipitated by a previous lot of this antibody (lot A301-737A-1). For blotting immunoprecipitated iASPP, A301-737A was used at 0.4 µg/ml. **Detection:** Chemiluminescence with an exposure time of 3 minutes.