

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


linkedin.com/company/szaboscandic in



#### **BRD4** Antibody

**Antigen Affinity Purified** 

Rabbit Polyclonal



Catalog No. A301-985A50 Uniprot ID O60885 Lot No. 9 GeneID 23476

**APPLICATIONS** WB, IP, IHC **SPECIES REACTIVITY** Human, Mouse

**AMOUNT** 50 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to BRD4 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A301-985A50 maps to a region between residue 1312 and 1362

of human bromodomain-containing protein 4 using the numbering given in entry

NP\_490597.1 (GeneID 23476).

Immunoglobulin concentration was determined using Beer's Law where 1 mg/mL IgG has an

A280 of 1.4.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:10,000 – 1:25,000 Immunoprecipitation 6 µg/mg lysate

Immunohistochemistry 1:1000 - 1:5000. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

IHC HUMAN CONTROLS Breast Carcinoma, Ovarian Carcinoma

IHC MOUSE CONTROLS Renal Cell Carcinoma

ADDITIONAL INFO https://www.fortislife.com/p/A301-985A50


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.


Michael Spencer, PhD

Date: May 14, 2024


BRD4 Antibody A301-985A50




Detection of human BRD4 by western blot. Samples: Whole cell lysate (10  $\mu$ g) from HeLa, HEK293T, Hep-G2, A-549, and SW620 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-BRD4 antibody (A301-985A50 lot 9) used for WB at 0.04  $\mu$ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.



**Detection of human BRD4 by immunohistochemistry.** *Sample:* FFPE section of prostate carcinoma. *Antibody:* Affinity purified rabbit anti-BRD4 antibody (A301-985A50 lot 9). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).




Detection of human BRD4 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg per IP; 5% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-BRD4 antibody (A301-985A50 lot 9) used for IP at 6 μg per reaction. BRD4 was also immunoprecipitated by a previous lot of this antibody (A301-985A50 lot 8), a recombinant monoclonal antibody [BL-151-6F11](A700-005) and an antibody against a different epitope of BRD4 [BL-149-2H5](A700-004A). For blotting immunoprecipitated BRD4, A301-985A was used at 0.04 μg/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.



Detection of mouse BRD4 by western blot. Samples: Whole cell lysate (5  $\mu$ g) from RenCa, TCMK-1, NIH 3T3, and CT26 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-BRD4 antibody (A301-985A50 lot 9) used for WB at 0.04  $\mu$ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

BRD4 Antibody A301-985A50



**Detection of mouse BRD4 by immunohistochemistry.** *Sample:* FFPE section of mouse renal cell carcinoma. *Antibody:* Affinity purified rabbit anti–BRD4 antibody (A301–985A50 lot 9). *Secondary:* HRP–conjugated goat anti–rabbit IgG (A120–501P).