

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



#### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# GSK3-beta Recombinant Monoclonal Antibody [BLR223K]



Rabbit Recombinant Monoclonal

Purified RefSeq ID NP\_002084.2

Catalog No. A700-223CF Uniprot ID P49841

Lot No. 1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

**AMOUNT** 100 μl

 $\textbf{CONCENTRATION} \qquad \quad 1000 \; \mu \text{g/ml}$ 

**STORAGE/SHELF LIFE**  $2 - 8^{\circ}\text{C} / 1$  year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Borate Buffered Saline (BBS) pH 8.2 with 0.09% Sodium Azide, BSA-Free

**ISOTYPE** IgG

CLONE # BLR223K
ORIGIN USA

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 370 and the C-terminus (residue 420) of human glycogen synthase kinase-3 beta using the numbering given in entry NP\_002084.2 (Gene ID 2932).

#### **APPLICATION NOTES**

All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent. Primary antibody is incubated overnight.

Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain Antibody (A120-101P).

Western blots of immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP Conjugate (A120-113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.

A700–223CF is the carrier–free formulation of A700–223. This product is optimized for conjugation with enzymes, fluorochromes, biotin, radioisotopes, oligonucleotides, microspheres, and other reagents. The optimal experimental concentration of the antibody post–conjugation must be determined by the investigator.

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com Orders: orders@fortislife.com • Support: technical@fortislife.com This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: April 19, 2024