



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# VE-Cadherin Recombinant Monoclonal Antibody [BLR239K]


Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_033998.2
Catalog No.	A700-239-T	Uniprot ID	P55284
Lot No.	A700-239-T-1	GenID	12562

---


<b>APPLICATIONS</b>	WB, IP, IHC, ICC
<b>SPECIES REACTIVITY</b>	Mouse
<b>AMOUNT</b>	10 µl (5+ tests)
<b>CONCENTRATION</b>	500 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 - 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>CLONE #</b>	BLR239K
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Recombinant antibody was purified from cell culture supernatant.  Immunogen was a recombinant protein representing the extracellular domain of mouse Cadherin-5 (aka VE-Cadherin, residues 25-595, NP_033998.2).
<b>APPLICATIONS</b>	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.  Western Blot                      1:1,000  Immunoprecipitation            12 µl/1 mg lysate  Immunohistochemistry        1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  Immunocytochemistry        1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.
<b>IHC MOUSE CONTROLS</b>	Gut, Lung, Uterus, bEnd Cells
<b>ADDITIONAL INFO</b>	<a href="https://www.fortislife.com/p/A700-239-T">https://www.fortislife.com/p/A700-239-T</a> Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: December 19, 2022




#### Detection of mouse VE-Cadherin by western blot.

**Samples:** Whole cell lysate (2 µg) prepared using NETN buffer from NIH 3T3, CT26, bEnd.3, TCMK-1, and BW5147.3 cells prepared using NETN lysis buffer. **Antibody:** Rabbit anti-VE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239-T lot 1) used at 1:1000. **Secondary:** HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).




**Detection of mouse VE-Cadherin by immunohistochemistry.** **Sample:** FFPE section of mouse lung. **Antibody:** Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239-T). **Secondary:** HRP-conjugated goat anti-rabbit IgG (A120-501P).



#### Detection of mouse VE-Cadherin by western blot of immunoprecipitates.

**Samples:** Whole cell lysate (1.0 mg per IP reaction; 5% of IP loaded) from bEnd.3 cells prepared using NETN lysis buffer. **Antibodies:** Rabbit anti-VE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239-T lot 1) used for IP at 12 µl/mg lysate. VE-Cadherin was also immunoprecipitated by a second antibody against a different epitope of VE-Cadherin (BL-2609B-4B7). For blotting immunoprecipitated VE-Cadherin, A700-239-T was used at 1:1000. Chemiluminescence with an exposure time of 1 second.



**Detection of mouse VE-Cadherin by immunohistochemistry.** **Sample:** FFPE section of mouse uterus. **Antibody:** Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239-T). **Secondary:** HRP-conjugated goat anti-rabbit IgG (A120-501P).



**Detection of mouse VE-Cadherin by immunocytochemistry.** *Sample:* FFPE section of bEnd cells. *Antibody:* Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239-T). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).