

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


VE-Cadherin Recombinant Monoclonal Antibody [BLR239K]




Rabbit Recombinant Monoclonal				
Purified			RefSeq ID	NP_033998.2
Catalog No.	No. A700-239		Uniprot ID	P55284
Lot No.	A700-2	239–1	GenelD	12562
APPLICATIONS		WB, IP, IHC, ICC		
SPECIES REACTIVITY		Mouse		
AMOUNT		100 µl (50+ tests)		
CONCENTRATION		500 μg/ml		
STORAGE/SHELF LIFE		2 - 8°C / 1 year from date of receipt		
PHYSICAL STATE		Liquid		
BUFFER		Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide		
ISOTYPE		IgG		
CLONE #		BLR239K		
ORIGIN		USA		
PRODUCTION PROCEDURES		Recombinant antibody was purified from cell culture supernatant.		
		Immunogen was a recombinant protein representing the extracellular domain of mouse Cadherin-5 (aka VE-Cadherin, residues 25-595, NP_033998.2).		
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.		
		Western Blot	1:	1,000
		Immunoprecipita	ation 12	µl/1 mg lysate
		Immunohistoche		100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is commended for FFPE tissue sections.
		Immunocytocher	,	100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is commended for FFPE cell sections.
IHC MOUSE CONTROLS Gut, Lung, Ute		Gut, Lung, Uterus	s, bEnd Cell	S
ADDITIONAL INFO		https://www.fortislife.com/p/A700-239 Use the link above to view SDS, a current list of citations, and other product specific information.		

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: December 19, 2022

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.




Detection of mouse VE–Cadherin by western blot. Samples: Whole cell lysate (2 µg) prepared using NETN buffer from NIH 3T3, CT26, bEnd.3, TCMK–1, and BW5147.3 cells prepared using NETN lysis buffer. Antibody: Rabbit anti–VE–Cadherin recombinant monoclonal antibody [BLR239K] (A700–239 lot 1) used at 1:1000. Secondary: HRP–conjugated goat anti–rabbit IgG (A120–101P). Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti–Actin recombinant monoclonal antibody [BLR057F] (A700–057).




Detection of mouse VE-Cadherin by

immunohistochemistry. *Sample:* FFPE section of mouse lung. *Antibody:* Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of mouse VE-Cadherin by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 5% of IP loaded) from bEnd.3 cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-VE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239 lot 1) used for IP at 12 μ l/mg lysate. VE-Cadherin was also immunoprecipitated by a second antibody against a different epitope of VE-Cadherin (BL-2609B-4B7). For blotting immunoprecipitated VE-Cadherin, A700-239 was used at 1:1000. Chemiluminescence with an exposure time of 1 second.



Detection of mouse VE-Cadherin by immunohistochemistry. *Sample:* FFPE section of mouse uterus. *Antibody:* Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. She information provided in this data sheet is believed to be material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.

3



Detection of mouse VE-Cadherin by immunocytochemistry. *Sample:* FFPE section of bEnd cells. *Antibody:* Rabbit anti-mVE-Cadherin recombinant monoclonal antibody [BLR239K] (A700-239). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).

Phone: 800.338.9579 · Fax: 866.597.6105 · Web: www.fortislife.com Orders: orders@fortislife.com · Support: technical@fortislife.com