

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



CD94 Recombinant Monoclonal Antibody [BLR264L]



Rabbit Recombinant Monoclonal

Purified RefSeq ID NP_002253.2

Catalog No. A700-264-T Uniprot ID Q13241 Lot No. 1 GeneID 3824

APPLICATIONS WB, IP SPECIES REACTIVITY Human

 AMOUNT
 20 μl (2 blots)

 CONCENTRATION
 250 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR264L
ORIGIN USA

PRODUCTION Recombinant antibody was purified from cell culture supernatant.

PROCEDURES

Immunogen was a recombinant protein representing the extracellular domain of human CD94 (residues 32–179, NP_002253.2).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000

Immunoprecipitation 20 µl/1 mg lysate


ADDITIONAL INFO https://www.fortislife.com/p/A700-264-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: May 31, 2023





Detection of human CD94 by western blot. Samples: Whole cell lysate (10 μg) from Jurkat, NK-92, and HEK293T cells prepared using NETN lysis buffer. Antibody: Rabbit anti-CD94 recombinant monoclonal antibody [BLR264L] (A700-264 lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Detection: Chemiluminescence with an exposure time of 3 minutes. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).

Detection of human CD94 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg per IP; 10% of IP loaded) from NK-92 cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-CD94 recombinant monoclonal antibody [BLR264L] (A700–264 lot 1) used for IP at 20 µl/mg lysate. CD94 was also immunoprecipitated by a second antibody against a different epitope of CD94 (BL-2778A-4D7). For blotting immunoprecipitated CD94, A700–264 was used at 1:1000. *Detection:* Chemiluminescence with an exposure time of 3 minutes.