



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# YTHDF2 Recombinant Monoclonal Antibody [BLR297M]


Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_057342.2
Catalog No.	A700-297	Uniprot ID	Q9Y5A9
Lot No.	1	GeneID	51441


---

<b>APPLICATIONS</b>	WB, Flow Cyt
<b>SPECIES REACTIVITY</b>	Human, Mouse
<b>AMOUNT</b>	100 µl (10 blots)
<b>CONCENTRATION</b>	500 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Phosphate Buffered Saline (PBS) with 0.1% BSA and 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>CLONE #</b>	BLR297M
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Recombinant antibody was purified from cell culture supernatant.  Immunogen was a peptide representing a region between residue 300 and 350 of human YTH domain-containing family protein 2 using the numbering given in entry NP_057342.2 (Gene ID 51441).
<b>APPLICATIONS</b>	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.  Western Blot                      1:1000 Flow Cytometry                  Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 x 10 <sup>6</sup> cells.
<b>APPLICATION NOTES</b>	All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent. Primary antibody is incubated overnight.  Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain Antibody (Cat. No. A120-101P).
<b>ADDITIONAL INFO</b>	<a href="https://www.fortislife.com/p/A700-297">https://www.fortislife.com/p/A700-297</a> Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: December 21, 2023




**Detection of human YTHDF2 by western blot.** *Samples:* Whole cell lysate (50  $\mu$ g) from CCRF-CEM, MOLT-4, K-562, Jurkat, and HEL 92.1.7 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-YTHDF2 recombinant monoclonal antibody [BLR297M] (A700-297 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). *Detection:* Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).



**Detection of human YTHDF2 (shaded) in Jurkat cells by flow cytometry.** *Antibody:* Rabbit anti-YTHDF2 recombinant monoclonal antibody [BLR297M] (A700-297) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).



**Detection of mouse YTHDF2 by western blot.** *Samples:* Whole cell lysate (25  $\mu$ g) from NIH 3T3, A20, and Ba/F3 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-YTHDF2 recombinant monoclonal antibody [BLR297M] (A700-297 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). *Detection:* Chemiluminescence with an exposure time of 10 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).



**Detection of mouse YTHDF2 (shaded) in NIH3T3 cells by flow cytometry.** *Antibody:* Rabbit anti-YTHDF2 recombinant monoclonal antibody [BLR297M] (A700-297) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).