

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



#### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# c-Rel Recombinant Monoclonal Antibody [BLR342M]



Rabbit Recombinant Monoclonal

Purified RefSeq ID NP\_002899.1 Catalog No. A700-342 Uniprot ID Q04864 Lot No. 1 GeneID 5966

**APPLICATIONS** WB, IP, IHC, ICC, Flow Cyt

SPECIES REACTIVITY Human

AMOUNT  $100 \mu l (50 + tests)$ 

CONCENTRATION 250 μg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Phosphate Buffered Saline (PBS) with 0.1% BSA and 0.09% Sodium Azide

**ISOTYPE** IgG

CLONE # BLR342M

ORIGIN USA

**PRODUCTION** 

Recombinant antibody was purified from cell culture supernatant.

**PROCEDURES** 

Immunogen was a peptide representing a region between residue 569 and the C-terminus

(residue 619) of human proto-oncogene c-Rel using the numbering given in entry

NP\_002899.1 (Gene ID 5966).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation 20 µl/mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE cell sections.

Flow Cytometry Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl

per 1 x  $10^6$  cells.

**APPLICATION NOTES** All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent.

Primary antibody is incubated overnight.

Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain

Antibody (A120-101P).

Western blots of immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP

Conjugate (A120-113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.

IHC HUMAN CONTROLS Appendix, Hodgkin Lymphoma, Ovarian Carcinoma, Prostate Carcinoma, Spleen, Tonsil, Daudi

Cells, MJ Cells, Raji Cells

ADDITIONAL INFO https://www.fortislife.com/p/A700-342

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer. PhD

Date: December 5, 2024