

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PAX8 Recombinant Monoclonal Antibody [BLR346N]





Rabbit Recombinant Monoclonal

Purified RefSeq ID NP_003457.1

Catalog No. A700-346 Uniprot ID Q06710 Lot No. 1 GeneID 7849

APPLICATIONS WB, IP, IHC, ICC

SPECIES REACTIVITY Human, Mouse

AMOUNT 100 µl (50+ tests)

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

1000 µg/ml

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) with 0.09% Sodium Azide, BSA-Free

ISOTYPE IgG

CLONE # BLR346N

ORIGIN USA

PRODUCTION Recombin

PROCEDURES

CONCENTRATION

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 225 and 275 of human Paired box protein Pax-8 using the numbering given in entry NP_003457.1 (Gene ID 7849).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation 6 µl/mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with Tris-EDTA pH 9.0 or pressure

cooker with citratebuffer pH 6.0 is recommended for FFPE tissue

sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with Tris-EDTA pH 9.0 or pressure

cooker with citratebuffer pH 6.0 is recommended for FFPE cell

sections

APPLICATION NOTES All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent.

Primary antibody is incubated overnight. Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain Antibody (A120-101P). Western blots of

immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP Conjugate (A120-

113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.

IHC HUMAN CONTROLS Fallopian Tube, Ovarian Carcinoma, OVCAR-3 Cells, OVCAR-4 Cells

IHC MOUSE CONTROLS Ovary, RenCa Cells

ADDITIONAL INFO https://www.fortislife.com/p/A700-346

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: January 6, 2025