

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



RUNX2 Recombinant Monoclonal Antibody [BLR360N]



Rabbit Recombinant Monoclonal

Purified RefSeg ID NP_001019801.3

Catalog No. A700-360 Uniprot ID Q13950 Lot No. 1 GeneID 860

APPLICATIONS WB, IP, IHC, ICC

SPECIES REACTIVITY Human, Mouse

AMOUNT 100 µl (50+ tests)

CONCENTRATION 500 μg/ml

STORAGE/SHELF LIFE $2 - 8^{\circ}\text{C} / 1$ year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR360N

ORIGIN USA

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 300 and 350 of human

Runt-related transcription factor 2 (RUNX2) using the numbering given in entry

NP_001019801.3 (Gene ID 8

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000

Immunoprecipitation 12 µl/mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE cell sections.

APPLICATION NOTES All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent.

Primary antibody is incubated overnight.

Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain

Antibody (A120-101P).

Western blots of immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP

Conjugate (A120-113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.

IHC HUMAN CONTROLS Hodgkin Lymphoma, Osteosarcoma, Tonsil, GaMG Cells, MDA-MB-231 Cells, SR Cells

IHC MOUSE CONTROLS C2C12 Cells, NIH 3T3 Cells

ADDITIONAL INFO https://www.fortislife.com/p/A700-360

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: March 28, 2025