



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic





Endoglin, Sf9, His Tag, human recombinant (rHuEndoglin-Sf9-His)

Catalog No: 94965
Lot No: XXXXX
Source: Sf9, Insect Cells
Synonyms: CD105, ENG, END, ORW, HHT1, ORW1, FLJ41744, Endoglin

Background

Endoglin is a type I membrane glycoprotein located on cell surfaces and is part of the TGF beta receptor complex. The protein consists of a homodimer of 180 kDa with disulfide links. It has been found on endothelial cells, activated macrophages, fibroblasts, and smooth muscle cells. Endoglin has been found to be part of the TGF-beta1 receptor complex. It thus may be involved in the binding of TGF-beta1, TGF-beta3, activin-A, BMP-2, and BMP-7. Beside TGF-beta signaling endoglin may have other functions. It has been postulated that endoglin is involved in the cytoskeletal organization affecting cell morphology and migration. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling. Its expression is regulated during heart development. Experimental mice without the endoglin gene die due to cardiovascular abnormalities.

Description

CD105 human recombinant extracellular domain produced in baculovirus is a homodimeric, glycosylated, polypeptide containing 586 amino acids and having a molecular mass of 61 kDa but as a result of glycosylation, migrates at 90 kDa under reducing conditions in SDS-PAGE. CD105 is fused to a C-terminal His Tag (6xHis) and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Endoglin was lyophilized from a concentrated (1 mg/ml) sterile solution containing 1xPBS.

Solubility

It is recommended to reconstitute the lyophilized CD105 in sterile PBS not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Endoglin, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CD105 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 95.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.



Amino Acid Sequence

MDRGTLPLAV ALLLASCSLS PTSLAETVHC DLQPGPERG EVTYTTSQVS KGCVAQAPNA ILEVHVLFLE FPTGPSQLEL TLQASKQNGT WPREVLLVLS VNSSVFHLQ ALGIPLHLAY NSSLVTFQEP PGVNNTTELPS FPKTQILEWA AERGPITSAA ELNDPQSILL RLGQAQGSLS FCMLEASQDM GRTLEWRPRT PALVRGCHE GVAGHKEAHI LRVLPGHSAG PRTVTVKVEL SCAPGDLDLAV LILQGPPYVS WLIDANHNMQ IWTTGEYSFK IFPEKNIRGF KLPDTPQGLL GEARMLNASI VASFVELPLA SIVSLHASSC GGRLQTSPAP IQTTPPKDTc SPELLMSLIQ TKCADDAMTL VLKKELVAHL KCTITGLTFW DPSCEAEDRG DKFVLRSAYS SCGMQVSASM ISNEAVVNIL SSSSPQRKKV HCLNMDLSF QLGLYLSPHF LQASNTIEPG QQSFSVQVRVS PSVSEFLQL DSCHLDDLGPE GGTVELIQGR AAKGNCSVSL SPSPSEGDPRF SFLLHFYTVP IPKTGTLSCt VALRPKTGS

Activity

Measured by its ability to bind with rhTGF-beta RII/Fc in a functional ELISA. Optimal dilutions should be determined by each laboratory for each application.

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.