

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Interferon alpha2b, human recombinant (rHuIFN-alpha2b-Yeast)

Catalog No: 94990 Lot No: XXXXX

Source: Saccharomyces cerevisiae

Synonyms: Interferon alpha 2b, IFNA, INFA2, IFN-? 2b, MGC125764, MGC125765

Background

IFN-alpha is produced by macrophages and has antiviral activities. Interferon stimulates the production of two enzymes: protein kinase and an oligoadenylate synthetase.

Description

Interferon alpha 2b human recombinant produced in yeast is a single, glycosylated, polypeptide chain containing 165 amino acids and having a molecular mass of approximately 19 kDa. The Interferon alpha 2b gene was obtained from human leukocytes. IFN-a 2b is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile filtered white lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered concentrated (1 mg/ml) solution in PBS, pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized glycosilated IFN-a 2b in sterile 18 M Ω -cm H $_2$ O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized glycosilated IFN-a 2b, although stable at room temperature for 3 weeks, should be stored desiccated below - 18°C. Upon reconstitution IFN-alpha 2b should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity

Greater than 98.0% as determined by (a) Analysis by RP-HPLC, (b) Analysis by SDS-PAGE.

Amino Acid Sequence

The sequence of the first five N-terminal amino acids was determined and was found to be Cys-Asp-Leu-Pro-Gln.

Activity

The specific activity as determined in a viral resistance assay was found to be no less than 300,000,000 IU/mg.

Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.