

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Activin-A, mouse recombinant (rmActivin A)

Catalog No: 97563 Lot No: XXXXX Source: E. coli

Synonyms: Inhba, Inhibin beta A, FSH releasing protein

Background

Activins are homodimers or heterodimers of the different ß subunit isoforms, part of the TGFß family. Mature Activin A has two 116 amino acids residues ßA subunits (ßA-ßA). Activin displays an extensive variety of biological activities, including mesoderm induction, neural cell differentiation, bone remodelling, haematopoiesis, and reproductive physiology. Activins takes part in the production and regulation of hormones such as FSH, LH, GnRH and ACTH. Cells that are identified to express Activin A include fibroblasts, endothelial cells, hepatocytes, vascular smooth muscle cells, macrophages, keratinocytes, osteoclasts, bone marrow monocytes, prostatic epithelium, neurons, chondrocytes, osteoblasts, Leydig cells, Sertoli cells, and ovarian granulosa cells.

Description

Active form Activin-A murine recombinant produced in *E. coli* is a homodimeric, non-glycosylated, polypeptide chain containing 2 x 117 amino acids and having a molecular weight of 26.2 kDa. Active form Activin-A is purified by standard chromatographic techniques.

Physical Appearance

Lyophilized freeze dried powder.

Formulation

Mouse Activin-A was lyophilized from a concentrated 1 mg/ml protein solution containing 0.02% TFA.

Solubility

INHBA protein should be reconstituted in distilled pyrogen free water to a concentration of 100 μ g/ml which can then be further diluted to other aqueous solutions.

Stability

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Repeated freezing and thawing is not recommended.

Purity

Greater than 95% as observed by SDS-PAGE.

Amino Acid Sequence

MGLECDGKVN ICCKKQFFVS FKDIGWNDWI IAPSGYHANY CEGECPSHIA GTSGSSLSFH STVINHYRMR GHSPFANLKS CCVPTKLRPM SMLYYDDGQN IIKKDIQNMI VEECGCS

Activity

Biological activity is assessed by the ability to induce cytoxicity of MPC-11 cells and was found to be 1 - 1.5 ng/ml corresponding to a specific activity of 666,667 - 1,000,000 units/mg.





Usage

This product is offered by Biomol for research purposes only. Not for diagnostic purposes or human use. It may not be resold or used to manufacture commercial products without written approval of Biomol GmbH.