

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



HER4 (ErbB4), Fc Fusion (IgG1), Avi-Tag, Biotin-Labeled Recombinant

Catalog: 102239 Lot: 250227

Product Information

Description: Recombinant human HER4 (human epidermal growth factor receptor 4), also known as

ErbB4, encompassing amino acids 26-651 (extracellular domain). This construct contains an Fc domain of human IgG1 fused to the C-terminus, followed by a C-terminal Avi-Tag $^{\text{TM}}$. The recombinant protein was enzymatically biotinylated using the Avi-Tag $^{\text{TM}}$

and affinity purified.

Background: HER4, also known as ErbB4 or receptor-tyrosine kinase 4, is a receptor tyrosine kinase

protein of the EGFR family. It is required for the development of the heart, nervous system, and mammary gland, and can also be found in the postsynaptic side of neuromuscular junctions and epidermis. HER4 is activated by neuroregulin 2-4, betacellulin and heparin binding EGF-like growth factor (HB-EGF) and is involved in mitogenesis and cell differentiation. HER4 undergoes a conformational change upon ligand binding and forms homodimers or heterodimers with other HER proteins to activate the MAPK (mitogen-activated protein kinases) and PI3K (phosphoinositide 3-kinase)/AKT pathways. Mutations in this protein can result in cancer but have also been linked to ALS (amyotrophic lateral sclerosis) type 19 and schizophrenia. The role of this protein as oncogene or tumor suppressor is still not fully determined, potentially being dependent on the cancer type. Several HER4 inhibitors have been developed and are undergoing clinical trials. Afatinib, a pan-HER inhibitor, has been approved for the treatment of NSCLC (non-small cell lung cancer), but may cause acute liver injury and death. The development of highly specific inhibitors for HER4 may prove beneficial in

oncology.

Species: Human

Construct: HER4 (26-651-Fc(IgG1)-Avi)-(Biotin)

Concentration:0.69 mg/mlExpression System:HEK293Purity:≥90%

Format: Aqueous buffer solution.

Formulated In: 8 mM phosphate, pH 7.4, 110 mM NaCl, 2.2 mM KCl, and 20% glycerol

MW: 98.7 kDa + glycans

Glycosylation: This protein runs at a higher MW by SDS-PAGE due to glycosylation.

Label: This protein is enzymatically biotinylated using Avi-Tag[™] technology. Biotinylation

confirmed to be ≥90%.

Genbank Accession: NM_005235.3

Stability: At least 6 months at -80°C.

Storage: -80°C

Instructions for Use: Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before

opening. Aliquot into small volumes and flash freeze for long term storage. Avoid

multiple freeze/thaw cycles.

Assay Conditions: The NRG1β Protein, Human (HEK293) was coated onto a 96-well plate overnight at 4°C

(25 ng/well). The plate was washed 3 times with PBST and blocked using 200 μ l of Blocking Buffer 3 (#79743) for 2 hours at Room Temperature (RT). After removing the blocking buffer, 50 μ l/well of purified HER4 (ErbB4), Fc Fusion (IgG1), Avi-Tag, Biotin-Labeled Recombinant (#102239), serially diluted in 1x PP-02 Buffer, was added for 1 hour at RT. After washing, the plate was incubated for 1 hour with HRP-labeled anti-FC antibody diluted 1:1000 in Blocking Buffer 3 (50 μ l/well) and washed again. Finally, 100 μ l of ECL substrate was added to each well and chemiluminescence was read

immediately.

Applications: Useful for binding studies.



HER4 (ErbB4), Fc Fusion (IgG1), Avi-Tag, Biotin-Labeled Recombinant


Catalog: 102239 Lot: 250227

Quality Control Data


4-20% SDS-PAGE Coomassie Staining

1 2 - 250 kDa - 150 kDa - 100 kDa - 75 kDa - 50 kDa - 37 kDa - 25 kDa - 20 kDa - 15 kDa - 10 kDa

Binding of HER4 (ErbB4) to NRG1β



Biotin-Avidin Pulldown



- 1. Beads
- 2. Flow Thru
- 3. Control
- 4. Standards

90 % biotin labeled.

* Avidin from beads.

