



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)




## Product Information

<b>Description:</b>	Recombinant human RNF43 (RING finger protein 43) encompassing amino acids 44-197. This construct contains a C-terminal Avi-tag™ followed by an His-tag (6xHis). This protein was affinity purified.
<b>Background:</b>	RNF43, or ring finger protein 43, is an E3 ubiquitin ligase that acts a tumor suppressor to negatively regulate the WNT (wingless-related integration site) signaling pathway through promoting the degradation of frizzled WNT receptors. Loss of function mutations in RNF43 have been shown to be a predictive biomarker in favorable response to anti-BRAF (B-Raf proto-oncogene serine/threonine kinase)/EGFR (epidermal growth factor receptor) therapies in colorectal cancer patients. RNF43 mutations are frequent predictors, and therefore of high clinical interest, in pancreatic, gastric, and endometrial cancers. Further studies will reveal more on the cancer signaling pathways and combinatory therapies involving RNF43.
<b>Species:</b>	Human
<b>Construct:</b>	RNF43 (44-197-Avi-His)
<b>Concentration:</b>	1.11 mg/ml
<b>Expression System:</b>	HEK293
<b>Purity:</b>	≥90%
<b>Format:</b>	Aqueous buffer solution.
<b>Formulated In:</b>	8 mM phosphate, 110 mM NaCl, 2.2 mM KCl, pH 7.4, and 20% glycerol.
<b>MW:</b>	20.2 kDa + glycans
<b>Glycosylation:</b>	This protein runs at a higher SDS-PAGE due to glycosylation.
<b>Genbank Accession:</b>	NM_017763.6
<b>Stability:</b>	At least 6 months at -80°C.
<b>Storage:</b>	-80°C
<b>Instructions for Use:</b>	Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.
<b>Assay Conditions:</b>	A 96-well plate was coated with RNF43, Avi-His-Tag Recombinant (100 ng/well in PBS) at 4°C overnight. The plate was washed three times with PBST, and 200 µl Blocking Buffer 3 was added for 2 hours at room temperature (RT). After three washes in PBST, a 1:2 serial dilution of RSPO2-Fc protein prepared in PP-02 Assay Buffer (BPS Bioscience #82620), was added and incubated at RT for 1 hour. The plate was washed three times and incubated with Anti-human Fc-HRP conjugate diluted of 1:1000 in Blocking Buffer 3 for 1 hour at RT. After three washes, 100 µl of the ELISA ECL Substrate (BPS Bioscience #79670) was added to each well and chemiluminescence was measured.
<b>Applications:</b>	Useful for binding assays.

Quality Control Data

4-20% SDS-PAGE Coomassie Staining



RNF43 Activity

