



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Product Information

Description:	Recombinant human EGFR (epidermal growth factor receptor), transcript variant 1, encompassing amino acids 25-645 of the extracellular domain. This construct contains a C-terminal His-tag (10xHis) followed by an Avi-Tag™. The recombinant protein was affinity purified.
Background:	EGFR (epidermal growth factor receptor) is the receptor for members of the EGF family and is a transmembrane glycoprotein that has tyrosine kinase activity. Binding of epidermal growth factor to EGFR induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation, differentiation, motility, and cell survival. Activation of EGFR triggers mitogenic signaling in gastrointestinal mucosa, and its expression is upregulated in colon cancers and most neoplasms. EGFR activates the ERK (extracellular signal-regulated kinase)-signaling pathway in normal gastric epithelial and colon cancer cell lines. Inactivation of EGFR with selective inhibitors significantly reduces ERK2 activation, c-fos mRNA expression and cell proliferation, opening new therapeutic opportunities.
Species:	Human
Construct:	EGFR (25-645-His-Avi)
Concentration:	1.49 mg/ml
Expression System:	HEK293
Purity:	≥90%
Format:	Aqueous buffer solution.
Formulated In:	8 mM phosphate, pH 7.4, 110 mM NaCl, 2.2 mM KCl, and 20% glycerol
MW:	73 kDa + glycans
Glycosylation:	This protein runs at a higher MW by SDS-PAGE due to glycosylation.
Genbank Accession:	NM_005228
Stability:	At least 6 months at -80°C.
Storage:	-80°C
Instructions for Use:	Thaw on ice and gently mix prior to use. DO NOT VORTEX. Perform a quick spin before opening. Aliquot into small volumes and flash freeze for long term storage. Avoid multiple freeze/thaw cycles.
Assay Conditions:	A white 96-well plate was precoated with Anti-EGFR, Avi-Tag Antibody (#102551) (100 ng/well). Blocking Buffer 3 (#79743) and PBST were used for blocking and washing steps, respectively. For protein titration a serial dilution of EGFR, His-Avi-Tag Recombinant (#102675) at the highest concentration of 4 ng/μl (200 ng/ reaction) was prepared in 1x PP-02 Assay Buffer. Diluted EGFR was then added to the wells and incubated at Room Temperature (RT) for 1 hour. The plate was then washed again and incubated with Anti-His-HRP diluted 1:1000 in Blocking Buffer 3 for 1 hour at RT. After the final wash, ELISA ECL substrate mix (#79670) was added to each well and chemiluminescence was read.
Applications:	Useful for binding assays.

Quality Control Data

