



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Information

Construct:	LRRK2 (G2019S) (GST-968-end)
Mutation:	G2019S
Concentration:	0.10 mg/ml
Species:	Human
Formulated In:	50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, 25% glycerol.
Expression System:	Sf9
Format:	Aqueous buffer solution
Stability:	At least 6 months at -80°C. Avoid freeze/thaw cycles.
Storage:	-80°C
Genbank Accession:	NM_198578
MW:	210 kDa
Purity:	70%
Specific Activity:	4.2 pmol/min/μg
Assay Conditions:	Kinase activity was measured using ADP-Glo™ Kinase Assay kit (Promega; Cat# V9101) which quantifies the amount of ADP produced by the LRRK2 (G2019S) reaction. The ADP-Glo™ Reagent is added to terminate the reaction and quench the remaining ATP. The Kinase Detection Reagent is then added to convert ADP to ATP and to measure the newly converted ATP using a luciferase/luciferin reaction. Assay: Kinase activity was measured using substrate LRRKtide (RLGRDKYKTLRQIRQ) (stock 1 mg/ml). Increasing amounts of LRRK2 (G2019S) were incubated with a final concentration of 0.2 mg/ml substrate in 40 mM Tris-HCl, pH 7.4, 20 mM MgCl₂, 0.1 mg/ml BSA, 50 μM fresh DTT, and 25 μM ATP. The total incubation time was 40 minutes at room temperature. The reaction was terminated by the addition of 5 μl of ADP-Glo™ Reagent and a subsequent incubation at room temperature for 40 minutes. Luminescence was measured by the addition of 10 μl Kinase Detection Reagent followed by incubation at room temperature for an additional 30 minutes. The blank was determined from a “no kinase” sample.
Applications:	Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Quality Control Data

4-20% SDS-Page Coomassie Staining



Specific Activity

