



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

### Description

Adeno-Associated Virus serotype 8 (AAV8) was isolated from rhesus monkey tissue, and the AAV8 rep and cap nucleotide sequences have 88% homology with AAV7 and 82% with AAV2. AAV8 exhibits greater transduction efficiency in the liver than other AAV serotypes. AAV8 and 9 have recently been used to correct disease-causing mutations and improve muscle function in mouse models of Duchenne muscular dystrophy.

These AAV particles constitutively express the firefly (*Photinus pyralis*) luciferase and mCherry genes connected via a T2A linker, under the control of a CMV promoter. The T2A self-cleaving peptide (derived from *Thosea asigna* virus 2A) leads to the efficient cleavage of the transcript and expression of luciferase and mCherry as two separate proteins.

### Application(s)

- Use as a positive control for transduction
- Optimize transduction assays and track protein expression over time

### Serotype


Wild-type AAV Serotype 8

### Formulation

AAV8 was produced in HEK293-AAV cells and is supplied in PBS-MK (PBS Magnesium-Potassium) buffer containing 0.01% Pluronic F68.

### Purification

The purity of the AAV particles was confirmed to be greater than 90% by staining with One-Step Lumitein™ UV Protein Gel Stain (Biotium #21005-1L). Purity will vary with each lot; the exact value will be provided with each shipment.



*Figure 1. Purified AAV8 Luciferase-mCherry particles.*

Staining of a 4-20% SDS-PAGE gel. The protein ladder is in lane 1, and  $2 \times 10^9$  GC (genome copy number) of AAV8 is shown in lane 2. Additional lanes between 1 and 2 were removed from the figure for clarity. AAV viral proteins VP1, VP2, and VP3 are labeled.

**Titer**


Two vials (50  $\mu$ l x 2) of AAV at a titer  $\geq 1 \times 10^{12}$  TU/ml. The titer is determined by qPCR and will vary with each lot; the exact value is provided with each shipment.

**Storage**

AAV is shipped with dry ice. For long-term storage, it is recommended to store AAV at  $-80^{\circ}\text{C}$ . Avoid repeated freeze-thaw cycles. Titers can drop significantly with each freeze-thaw cycle.


**Biosafety**

Recombinant AAV is inherently replication-deficient and not known to cause any human diseases. Additionally, following transduction, AAV vectors exist episomally and do not integrate into or disrupt the host cell's genome. AAV requires the use of a Biosafety Level 1 facility. BPS Bioscience recommends following all local, federal, state, and institutional regulations and using all appropriate safety precautions.

**Validation Data**

*Figure 2. Transduction of HEK293 cells using AAV8 Luciferase-mCherry particles.*

$1 \times 10^5$  cells/well were transduced in a 6-well plate with AAV8 Luciferase-mCherry at an MOI of  $2 \times 10^4$ . After 72 hours of transduction, mCherry expression in the target cells was observed under a fluorescence microscope. mCherry expression was stable over time and still observed 30 days after transduction.



*Figure 3. Luciferase activity of HEK293 cells transduced by AAV8 Luciferase-mCherry particles. 1 x 10<sup>5</sup> cells/well were transduced in a 6-well plate with AAV8 Luciferase-mCherry at an MOI of 2 x 10<sup>4</sup>. After 72 hours of transduction, transduced cells or parental HEK293 cells were seeded in a 96-well plate at a density of 2 x 10<sup>4</sup> cells/well, and luciferase activity was measured using the ONE-Step™ luciferase assay system (BPS Bioscience #60690)*

### Troubleshooting Guide

Visit [bpsbioscience.com/lentivirus-faq](https://bpsbioscience.com/lentivirus-faq) for detailed troubleshooting instructions. For all further questions, please email [support@bpsbioscience.com](mailto:support@bpsbioscience.com).

### Related Products

<i>Products</i>	<i>Catalog #</i>	<i>Size</i>
AAV1 ZsGreen	78443	50 µl x 2
AAV2 ZsGreen	78444	50 µl x 2
AAV5 ZsGreen	78447	50 µl x 2
AAV8 ZsGreen	78449	50 µl x 2
AAV1 Luciferase-mCherry	78470	50 µl x 2
AAV6 Luciferase-mCherry	78475	50 µl x 2
AAV9 Luciferase-mCherry	78477	50 µl x 2
AAV2 Luciferase-eGFP	78462	50 µl x 2
AAV8 Luciferase-eGFP	78467	50 µl x 2
AAV9 Luciferase-eGFP	78468	50 µl x 2