

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Description

Adeno-Associated Virus Serotype 1 (AAV1) exhibits high homology with other AAV serotypes. AAV1 efficiently transduces muscle tissue, as determined by a region of the capsid protein VP1 (amino acids 350 to 430) which functions as a major determinant of tissue tropism.

Cas9 is an endonuclease enzyme that introduces a double stranded break into the DNA when recruited to a specific DNA sequence by the sgRNA (single guide RNA). This double stranded break is repaired in mammalian cells either through Non-Homologous End Joining or Homologous Recombination. Non-Homologous End Joining often results in the deletion or insertion of several base pairs at the cut site, which, when resulting in a frameshift, causes the functional inactivation of the gene.

SaCas9 (*Staphylococcus aureus* CRISPR-associated protein 9) has demonstrated high cutting efficiency in mammalian cells, and its smaller size makes it ideal for packaging into AAV. SaCas9 recognizes a longer protospacer adjacent motif (PAM) site, 5'-NNGRRT-3', than the more traditional SpCas9 (*Streptococcus pyogenes* CRISPR-associated protein 9). These AAV particles constitutively express SaCas9 under the control of a CMV promoter.

Application(s)

Express SaCas9 to enable genome editing

Serotype

Wild-type AAV Serotype 1

Formulation

AAV was produced in HEK293-AAV cells and is supplied in PBS-MK (PBS Magnesium-Potassium) buffer containing 0.01% Pluronic F68.

Purification

The purity of the AAV particles was confirmed to be greater than 90% by staining with One-Step Lumitein[™] UV Protein Gel Stain (Biotium #21005-1L). The purity will vary with each lot; the exact value will be provided with each shipment.




Figure 1. Purified AAV1 SaCas9 particles.

Staining of a 4-20% SDS-PAGE gel. The protein ladder is in lane 1, and 2 x 10⁹ GC (genome copy number) of AAV is shown in lane 2. AAV viral proteins VP1, VP2, and VP3 are labelled.



Titer

Two vials (50 μ l x 2) of AAV at a titer $\ge 1 \times 10^{12}$ TU/ml. The titer is determined by qPCR and will vary with each lot; the exact value will be provided with each shipment.

Storage

AAV is shipped with dry ice. For long-term storage, it is recommended to store AAV at -80°C. Avoid repeated freeze-thaw cycles. Titers can drop significantly with each freeze-thaw cycle.

Biosafety



Recombinant AAV is inherently replication-deficient and not known to cause any human diseases. Additionally, following transduction, AAV vectors exist episomally and do not integrate into or disrupt the host cell's genome. AAV requires the use of a Biosafety Level 1 facility. BPS Bioscience recommends following all local, federal, state, and institutional regulations and using all appropriate safety precautions.

Validation Data



Figure 2. Transduction of HEK293 cells using AAV1 SaCas9.

 1×10^5 cells were transduced in a 6-well plate with AAV1 SaCas9 at an MOI of 1×10^4 . After 72 hours of transduction, SaCas9 expression in the target cells was measured by Western blot using Anti-HA.11 Epitope Tag Antibody (Biolegend #901516).

Troubleshooting Guide

Visit bpsbioscience.com/lentivirus-faq for detailed troubleshooting instructions. For all further questions, please email support@bpsbioscience.com.

Catalog #	Size
78443	50 μl x 2
78449	50 μl x 2
78450	50 μl x 2
78452	50 μl x 2
78453	50 μl x 2
78461	50 μl x 2
78463	50 μl x 2
78475	50 μl x 2
78476	50 μl x 2
78480	50 μl x 2
	78443 78449 78450 78452 78453 78461 78463 78475 78476



2