



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Description

The eGFP/Firefly Luciferase RS4;11 Cell Line are engineered human lymphoblast RS4;11 cells expressing firefly luciferase and enhanced GFP (eGFP) driven by an EF1a promoter. The cells were generated by transduction with Firefly Luciferase-eGFP Lentivirus (BPS Bioscience #78741), which is a SIN (self-inactivating) lentivirus.

Background

RS4;11 is a lymphoblast cell line derived from the marrow of a 32-year-old white female patient with acute lymphoblastic leukemia (ALL). This cell line has a KMT2A (histone-lysine N-methyltransferase 2A) rearrangement, which despite its low frequency leads to acute leukemia. It expresses HLA-C alleles, that bind the most expressed KIRs (killer-cell immunoglobulin-like receptors) and it is resistant to NK attack. Its characteristics make it an ALL-cell model and an appropriate negative control in NK cytotoxicity assays. The presence of eGFP and luciferase allows for easy assay readouts, making this cell line a convenient choice.

Application

- *In vitro* and *in vivo* bioluminescence imaging (BLI) and fluorescence imaging.
- Use as negative control in NK cytotoxicity assays.

Materials Provided

Components	Format
2 vials of frozen cells	Each vial contains $>1 \times 10^6$ cells in 1 ml of Cell Freezing Medium (BPS Bioscience #79796)

Parental Cell Line

RS4;11, human lymphoblast cell line, suspension.

Mycoplasma Testing

The cell line has been screened to confirm the absence of Mycoplasma species.

Materials Required but Not Supplied

These materials are not supplied with the cell line but are necessary for cell culture and cellular assays. BPS Bioscience's reagents are validated and optimized for use with this cell line and are highly recommended for best results. Media components are provided in the Media Formulations section below.

Media Required for Cell Culture

Name	Ordering Information
Thaw Medium 2	BPS Bioscience #60184
Growth Medium 2D	BPS Bioscience #79639

Storage Conditions

Cells are shipped in dry ice and should immediately be thawed or stored in liquid nitrogen upon receipt. Do not use a -80°C freezer for long term storage. Contact technical support at support@bpsbioscience.com if the cells are not frozen in dry ice upon arrival.

Media Formulations

For best results, the use of validated and optimized media from BPS Bioscience is *highly recommended*. Other preparations or formulations of media may result in suboptimal performance.



Note: Thaw Media do *not* contain selective antibiotics. However, Growth Media *do* contain selective antibiotics, which are used to maintain selective pressure on the cell population expressing the gene of interest.

Cells should be grown at 37°C with 5% CO₂. BPS Bioscience's cell lines are stable for at least 10 passages when grown under proper conditions.

Media Required for Cell Culture

Thaw Medium 2 (BPS Bioscience #60184):

RPMI-1640 medium supplemented with 10% FBS, 1% Penicillin/Streptomycin.

Growth Medium 2D (BPS Bioscience #79639):

RPMI-1640 medium supplemented with 10% FBS, 1% Penicillin/Streptomycin plus 200 µg/ml of Hygromycin.

Cell Culture Protocol

Cell Thawing

1. Swirl the vial of frozen cells for approximately 60 seconds in a 37°C water bath. As soon as the cells are thawed (it may be slightly faster or slower than 60 seconds), quickly transfer the entire content of the vial to a tube containing 10 ml of pre-warmed Thaw Medium 2.

Note: Leaving the cells in the water bath at 37°C for too long will result in rapid loss of viability.

2. Immediately spin down the cells at 300 x g for 5 minutes, remove the medium and resuspend the cells in 5 ml of pre-warmed Thaw Medium 2.
3. Transfer the resuspended cells to a T25 flask or T75 flask and incubate at 37°C in a 5% CO₂ incubator.
4. After 24 hours of culture, check for cell viability. For a T25 flask, add 3-4 ml of Thaw Medium 2, and continue growing in a 5% CO₂ incubator at 37°C until the cells are ready to passage.
5. Cells should be passaged before they reach a density of 2 x 10⁶ cells/ml. Switch to Growth Medium 2D at first and subsequent passages.

Cell Passage

Dilute the cell suspension into new culture vessels before they reach a density of 2 x 10⁶ cells/ml, with Growth Medium 2D. The sub-cultivation ratio should maintain the cells between 0.2 x 10⁶ cells/ml and 2 x 10⁶ cells/ml.

Cell Freezing

1. Spin down the cells at 300 x g for 5 minutes. Remove the medium and resuspend the cell pellet in 4°C Cell Freezing Medium (BPS Bioscience #79796) at a density of ~2 x 10⁶ cells/ml.
2. Dispense 1 ml of cell suspension into each cryogenic vial. Place the vials in an insulated container for slow cooling and store at -80°C overnight.

Transfer the vials to liquid nitrogen the next day for long term storage.



Note: It is recommended to expand the cells and freeze at least 10 vials at an early passage for future use.

Validation Data




Figure 1. eGFP expression and luciferase activity in the eGFP/Firefly Luciferase RS4;11 Cell Line.

A. Luciferase activity in eGFP/Firefly Luciferase RS4;11 cells and RS4;11 parental cells was measured using One Step™ Luciferase Assay System (BPS Bioscience #60690). **B.** 25,000 eGFP/Firefly Luciferase RS4;11 cells and parental RS4;11 cells were analyzed by flow cytometry for eGFP expression. The RS4;11 Parental cells are shown in blue, and the eGFP/Firefly Luciferase RS4;11 cells are shown in green.




Figure 2. Luciferase activity-based NK cytotoxicity using eGFP/Luciferase K562 and eGFP/Luciferase RS4;11 as target cells.

NK cells were enriched and expanded from PBMCs using NK Expansion Kit (BPS Bioscience #78927) for 14 days. Expanded NK cells were co-cultured with eGFP/Luciferase K562 target cells (red; BPS Bioscience #78911) and eGFP/Luciferase RS4;11 control cells (blue) at different Effector: Target (E:T) cell ratios at 37°C for 4 hours. Luciferase activity was detected using ONE-Step™ Luciferase. Cytotoxicity potency (specific lysis) correlates with a decrease in luciferase signal in target cells. Luminescence results were used to generate killing curves.

Data shown is representative. For lot-specific information, please contact BPS Bioscience, Inc. at support@bpsbioscience.com.

License Disclosure

Visit bpsbioscience.com/license for the label license and other key information about this product.

Troubleshooting Guide

Visit bpsbioscience.com/cell-line-faq for detailed troubleshooting instructions. For all further questions, please email support@bpsbioscience.com.

Related Products

Products	Catalog #	Size
Firefly Luciferase Lentivirus	79692	500 µl x 2
Firefly Luciferase-eGFP Lentivirus	78741	500 µl x 2
eGFP/Firefly Luciferase U-87 MG Cell Line	78904	2 vials
eGFP/ Firefly Luciferase MM.1S Cell Line	78376	2 vials
eGFP/ Firefly Luciferase K562 Cell Line	78911	2 vials

Version 103123