



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6042 Cornerstone Court West, Suite B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet

STAT3 Reporter (Luc) - HEK293 Cell line (Puromycin)

Catalog #: 79800-P

Product Description

The STAT3 Reporter (Luc)-HEK293 cell line is designed for monitoring STAT3 signal transduction pathway. It contains a firefly luciferase gene driven by STAT3 response elements located upstream of the minimal TATA promoter. After activation by cytokines and growth factors, endogenous STAT3 binds to the DNA response elements, inducing transcription of the luciferase reporter gene.

Format

Each vial contains ~2 X 10⁶ cells in 1 ml of 10% DMSO.

Applications

- Monitor the STAT3 signaling pathway activity
- Screen for activators or inhibitors of the STAT3 signaling pathway

General Culture Conditions

Thaw Medium 1 (BPS Bioscience #60187): MEM medium (Hyclone #SH30024.01) supplemented with 10% FBS, 1% non-essential amino acids (Hyclone #SH30238.01), 1 mM Na pyruvate (Hyclone #SH30239.01), 1% Penicillin/Streptomycin (Hyclone #SV30010.01).

Growth Medium 1N (BPS Bioscience #79801): Thaw Medium 1 (BPS Bioscience #60187) and 0.5 µg/ml of Puromycin (InvivoGen, #ant-pr-1).

Cells should be grown at 37°C with 5% CO₂ using Growth Medium 1N.

Storage

Immediately upon receipt, store in liquid nitrogen.

To thaw the cells, it is recommended to quickly thaw the frozen cells from liquid nitrogen in a 37°C water-bath, and transfer to a tube containing 10 ml of Thaw Medium 1 (**no puromycin**). Spin down cells, resuspend cells in pre-warmed Thaw Medium 1 (**no puromycin**), transfer resuspended cells to a T25 flask and culture in 37°C CO₂ incubator. At first passage switch to Growth Medium 1N (**contains puromycin**). Cells should be split before they reach complete confluence.

To passage the cells, rinse cells with phosphate buffered saline (PBS), detach cells from culture vessel with Trypsin/EDTA, add Growth Medium 1N and transfer to a tube. Spin down cells, resuspend cells and seed appropriate aliquots of cell suspension into new culture vessels. Subcultivation ration: 1:5 to 1:10 weekly or twice a week.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court West, Suite B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Note: Just after thawing and at low density, the cells may grow at a slower rate. It is recommended to split the cells with ~ 1:4 ratio at the beginning of culturing. After several passages, the cell growth rate increases and the cells can be split with a higher ratio.

To freeze down the cells, rinse cells with phosphate buffered saline (PBS), and detach cells from culture vessel with Trypsin/EDTA. Add Growth Medium 1N and transfer to a tube, spin down cells, and resuspend in freezing medium (10% DMSO + 90% FBS) at $\sim 2 \times 10^6$ cells/ml. Dispense 1 ml of cell aliquots into cryogenic vials. Place vials in an insulated container for slow cooling and store at -80°C overnight. Transfer to liquid nitrogen the next day for storage. It is recommended to expand the cells and freeze down more than 10 vials of cells for future use at early passage.

Mycoplasma testing

The cell line has been screened using the PCR-based Venor™GeM Mycoplasma Detection kit (Sigma-Aldrich, #MP0025) to confirm the absence of *Mycoplasma* species.

Assay performance

The following assays are designed for 96-well format. To perform the assay in different tissue culture formats, the cell number and reagent volume should be scaled appropriately.

Materials Required but Not Supplied

- Human IL-6 (R&D Systems #206-IL)
- JAK inhibitor CP690550 (Cayman #11598)
- Anti-IL-6R antibody (R&D Systems #MAB227)
- Assay Medium: Thaw Medium 1 (BPS Bioscience #60187)
- Growth Medium 1N (BPS Bioscience #79801)
- 96-well tissue culture treated white clear-bottom assay plate (Corning #3610)
- One-Step luciferase assay system (BPS Bioscience #60690) or other luciferase reagents for measuring firefly luciferase activity
- Luminometer

A. Human IL-6 dose response

1. Harvest STAT3 reporter (Luc)-HEK293 cells and seed cells at a density of 30,000-40,000 cells per well into white opaque 96-well microplate in 90 μl of assay medium. Incubate cells at 37°C with 5% CO_2 overnight.
2. Add threefold serial dilution of IL-6 in 10 μl of assay medium to IL-6-stimulated wells.
 - a. Add 10 μl of assay medium to the unstimulated control wells (for measuring uninduced level of STAT3 reporter activity).
 - b. Add 100 μl of assay medium to cell-free control wells (for determining background luminescence).

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**


Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com

3. Incubate at 37°C with 5% CO₂ for 5-18 hours.
4. Prepare ONE-Step™ Luciferase Assay reagent as directed and add 100 µl per well. Incubate at room temperature for ~15 to 30 minutes and measure luminescence using a luminometer. Subtract background luminescence value from all measurements.

Figure 1. IL-6 Dose Response in STAT3 (Luc) Reporter HEK293 Cells. The results are shown as fold induction of luciferase reporter expression. Fold induction was determined by comparing values against the mean value for control cells without IL-6 treatment.

The EC₅₀ of IL-6 in this cell line is ~2 ng/ml.



B. Inhibition of IL-6 induced STAT3 activity by JAK inhibitor

1. Harvest STAT3 reporter (Luc)-HEK293 cells and seed cells at a density of 30,000-40,000 cells per well into white opaque 96-well microplate in 90 µl of assay medium. Incubate cells at 37°C with 5% CO₂ overnight.
2. Next day, treat cells with three-fold serial dilution of CP690550 (JAK inhibitor; Cayman, #11598) in 90 µl of assay medium. Incubate cells at 37°C with 5% CO₂ for 1-2 hours. For control cells without CP690550, change to 90 µl of assay medium with no treatment.
3. Set up each treatment in at least triplicates:
 - a. Add 10 µl of diluted human IL-6 in assay medium to stimulated wells (final IL-6 concentration = 10 ng/ml).
 - b. Add 10 µl of assay medium to the unstimulated control wells (for determining the basal activity).

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.


To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com

- c. Add 100 μ l of assay medium to cell-free control wells (for determining background luminescence).
4. Incubate at 37°C with 5% CO₂ for 5-18 hours.
5. Prepare ONE-Step™ Luciferase Assay reagent as directed and add 100 μ l per well. Incubate at room temperature for ~15 to 30 minutes and measure luminescence using a luminometer. Subtract background luminescence value from all measurements.
6. Data Analysis: Obtain background-subtracted luminescence by subtracting the average background luminescence (cell-free control wells) from the luminescence reading of all wells.

Figure 2. Inhibition of IL-6-induced Reporter Activity by JAK Inhibitor CP690550 in STAT3 (Luc) Reporter HEK293 Cells. The results are shown as percentage of luminescence. The background-subtracted luminescence of cells stimulated with IL-6 in the absence of CP690550 was set at 100%.



C. Inhibition of IL-6 induced STAT3 activity by anti-IL-6R antibody

1. Harvest STAT3 reporter (Luc)-HEK293 cells and seed cells at a density of 30,000 cells per well into white opaque 96-well microplate in 90 μ l of assay medium. Incubate cells at 37°C with 5% CO₂ overnight.
2. Next day, treat cells with three-fold serial dilution of anti-IL-6R antibody (R&D Systems, #MAB227) in 90 μ l of assay medium for one hour. For control cells without antibody treatment, change to 90 μ l of fresh assay medium.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com


Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court West, Suite B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

3. Set up each treatment in at least triplicates:
 - a. Add 10 μ l of diluted human IL-6 in assay medium to stimulated wells (final IL-6 concentration = 10 ng/ml).
 - b. Add 10 μ l of assay medium to the unstimulated control wells (for determining the basal activity).
 - c. Add 100 μ l of assay medium to cell-free control wells (for determining background luminescence).
4. Incubate at 37°C with 5% CO₂ for 18-24 hours.
5. Prepare ONE-Step™ Luciferase Assay reagent as directed and add 100 μ l per well. Incubate at room temperature for ~15 to 30 minutes and measure luminescence using a luminometer. Subtract background luminescence value from all measurements.
6. Data Analysis: Obtain background-subtracted luminescence by subtracting the average background luminescence (cell-free control wells) from the luminescence reading of all wells.

Figure 3. Inhibition of IL-6-induced Reporter Activity by Anti-IL-6R Antibody in STAT3 (Luc) Reporter HEK293 Cells. The results are shown as percentage of luminescence. The background-subtracted luminescence of cells stimulated with IL-6 in the absence of anti-IL-6R antibody was set at 100%.



References

1. Tian S., *et al.*, *Blood*. 1994; **84(6)**:1760-1764.
2. Zhong, Z., *et al.*, *Science*. 1994; **264(5155)**:95-98.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6042 Cornerstone Court West, Suite B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

License Disclosure

Purchase of this cell line grants you with a 10-year license to use this cell line in your immediate laboratory, for research use only. This license does not permit you to share, distribute, sell, sublicense, or otherwise make the cell line available for use to other laboratories, departments, research institutions, hospitals, universities, or biotech companies. The license does not permit use of this cell line in humans or for therapeutic or drug use. The license does not permit modification of the cell line in any way. Inappropriate use or distribution of this cell line will result in revocation of the license and result in an immediate cease of sales and distribution of BPS products to your laboratory. BPS does not warrant the suitability of the cell line for any particular use, and does not accept any liability in connection with the handling or use of the cell line. Modifications of this cell line, transfer to another facility, or commercial use of the cells may require a separate license and additional fees; contact sales@bpsbioscience.com for details. Publications using this cell line should reference BPS Bioscience, Inc., San Diego.

Related Products

Product	Cat. #	Size
Thaw Medium 1	60187	100 ml
Growth Medium 1N	79801	500 ml
STAT3 Reporter Kit	79730	500 rxns.
STAT3 Luciferase Reporter Lentivirus	79744	2 x 500 µl
Human IL-6	90196-B	20 µg
ONE-Step Luciferase Detection Reagent	60690-1	10 ml
STAT3, GST-tag	75003	20 µg
STAT5 Reporter (Luc) – Ba/F3 Cell line	79722	2 vials

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com