

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

Description

Firefly Luciferase A549 Cell Line is a lung cancer A549 cell line that expresses the firefly luciferase reporter driven by an EF1a promoter. This cell line has been generated by transduction with Firefly Luciferase Lentivirus (BPS Bioscience #78740-H).

This cell line has been validated in co-culture assays.

Background

A549 cells were isolated from the lung tissue of a lung cancer patient. This cell line is the most commonly used human non-small cell lung cancer (NSCLC) cell line for both basic cancer research and drug discovery. Firefly luciferase has been used as a sensitive reporter to study a wide range of biological responses. It was first cloned from the North American *Photinus pyralis* and catalyzes the oxidation of D-luciferin, in the presence of ATP and magnesium, emitting yellow light. This reaction has a high quantum yield, and both luciferase and luciferin have low toxicity. These characteristics contributed to Firefly luciferase becoming a commonly used tool. The use of firefly luciferase as reporter allows for easy readouts.

Application

- Use as target cells in CAR-T or CAR-NK co-culture killing assays.
- In vitro and in vivo bioluminescence imaging.

Materials Provided

Components	Format
2 vials of frozen cells	Each vial contains ≥1 x 10 ⁶ cells in 1 ml of Cell Freezing
	Medium (BPS Bioscience #79796)

Parental Cell Line

A549 is a human lung alveolar vessel carcinoma cell line. Adherent epithelial cells.

Mycoplasma Testing

The cell line has been screened to confirm the absence of Mycoplasma species.

Materials Required but Not Supplied



These materials are not supplied with the cell line but are necessary for cell culture and cellular assays. BPS Bioscience's reagents are validated and optimized for use with this cell line and are highly recommended for best results. Media components are provided in the Media Formulations section below.

Media Required for Cell Culture

Name	Ordering Information
Thaw Medium 6	BPS Bioscience #60183
Growth Medium 6E	BPS Bioscience #78385

Storage Conditions



Cells are shipped in dry ice and should immediately be thawed or stored in liquid nitrogen upon receipt. Do not use a -80°C freezer for long term storage. Contact technical support at support@bpsbioscience.com if the cells are not frozen in dry ice upon arrival.



Media Formulations

For best results, the use of validated and optimized media from BPS Bioscience is highly recommended. Other preparations or formulations of media may result in suboptimal performance.



Note: Thaw Media do *not* contain selective antibiotics. However, Growth Media *do* contain selective antibiotics, which are used to maintain selective pressure on the cell population expressing the gene of interest.

Cells should be grown at 37° C with 5% CO₂. BPS Bioscience's cell lines are stable for at least 10 passages when grown under proper conditions.

Media Required for Cell Culture

Thaw Medium 6 (BPS Bioscience #60183):

DMEM medium supplemented with 10% FBS, 1% Penicillin/Streptomycin.

Growth Medium 6E (BPS Bioscience #78385):

DMEM medium supplemented with 10% FBS, 1% Penicillin/Streptomycin plus 600 µg/ml Hygromycin B.

Cell Culture Protocol

Note: Note: A549 cells are derived from human material and thus the use of adequate safety precautions is recommended.

Cell Thawing

1. Swirl the vial of frozen cells for approximately 60 seconds in a 37°C water bath. As soon as the cells are thawed (it may be slightly faster or slower than 60 seconds), quickly transfer the entire contents of the vial to a tube containing 10 ml of pre-warmed Thaw Medium 6.

Note: Leaving the cells in the water bath at 37°C for too long will result in rapid loss of viability.

- 2. Immediately spin down the cells at 300 *x g* for 5 minutes, remove the medium and resuspend the cells in 5 ml of pre-warmed Thaw Medium 6.
- 3. Transfer the resuspended cells to a T25 flask and incubate at 37°C in a 5% CO2 incubator.
- 4. After 48-72 hours of culture, check for cell viability, change to fresh Thaw Medium 6, and continue growing in a 5% CO₂ incubator at 37°C until the cells are ready to passage.
- 5. Cells should be passaged before they reach 100% confluency. Switch to Growth Medium 6E for passage.

Cell Passage

- 1. Aspirate the medium, wash the cells with phosphate buffered saline (PBS) without Ca²⁺/Mg²⁺, and detach the cells from the culture vessel with 0.25% Trypsin/EDTA following volumes recommended for the cell vessel being used.
- 2. Once the cells have detached, add Growth Medium 6E and transfer to a tube.



- 3. Spin down cells at 300 *x g* for 5 minutes, remove the medium and resuspend the cells in Growth Medium 6E.
- 4. Seed into new culture vessels at the recommended sub-cultivation ratio of 1:4 to 1:5 weekly or twice per week.

Cell Freezing

- 1. Aspirate the medium, wash the cells with PBS without Ca²⁺/Mg²⁺, and detach the cells from the culture vessel with 0.25% Trypsin/EDTA following volumes recommended for the cell vessel being used.
- 2. Once the cells have detached, add Growth Medium 6E and transfer to a tube.
- 3. Spin down cells at 300 x g for 5 minutes.
- 4. Remove the medium and resuspend the cell pellet in 4°C Cell Freezing Medium (#79796) at a density of 2 x 10 cells/ml.
- 5. Dispense 1 ml of cell suspension into each cryogenic vial. Place the vials in an insulated container for slow cooling and store at -80°C overnight.
- 6. Transfer the vials to liquid nitrogen the next day for long term storage.



Note: It is recommended to expand the cells and freeze at least 10 vials at an early passage for future use.

Validation Data

Cytotoxicity assay of Anti-GPC3 CAR-T Cells using Firefly Luciferase A549 Cell Line as the target cells

This cell line was validated in a co-culture cytotoxicity assay. For details about the exact protocol used to evaluate the cytotoxicity function of Anti-GPC3 CAR-T Cells, refer to the anti-GPC3 CAR-T Cells datasheet (#82494).






Figure 1. Luciferase-based cytotoxicity assay of Anti-GPC3 CAR-T Cells using Firefly Luciferase A549 Cell Line as the target cells.

Anti-GPC3 CAR-T effector cells (#82494) were thawed and co-cultured with Firefly Luciferase A549 cells as the target cells for 24 hours at the indicated E:T ratios. The assay was performed in parallel with untransduced T cells as a negative control. The lysis of the target cells after 24 hours was determined by measuring luciferase activity with ONE-Step™ Assay System (#60690). The average background luminescence was subtracted from the luminescence reading of all wells. Results are expressed as Percent of Luminescence, with "Firefly Luciferase HepG2 cells" alone set to 100%. The luminescence signal is proportional to the amount of living cells remaining in the co-culture population.

Data is representative.

License Disclosure

Visit bpsbioscience.com/license for the label license and other key information about this product.

Troubleshooting Guide

Visit bpsbioscience.com/cell-line-faq for detailed troubleshooting instructions. For lot-specific information and all other questions, please email visit https://bpsbioscience.com/contact.

Related Products

_Products	Catalog #	Size
Firefly Luciferase Lentivirus (G418, Hygromycin and Puromycin)	79692	500 μl x 2
Firefly Luciferase Lentivirus EF1A Promoter/ G418, Hygromycin, or Puromycin)	78740	500 μl x 2
Firefly Luciferase-eGFP Lentivirus (G418 or Puromycin)	79980	500 μl x 2
Firefly Luciferase-eGFP Lentivirus (EF1A Promoter/ G418, Hygromycin, or Puromycin)	78741	500 μl x 2

Version 060925

