



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Description

Firefly Luciferase A375 Cell Line is an A375 human melanoma cell line engineered to express Firefly luciferase driven by a CMV promoter. This cell line was generated by transduction with Firefly Luciferase Lentivirus (#79692-P).

Background

A375 is a melanoma cell line isolated from a patient with malignant melanoma, that is used extensively in toxicology and oncology studies. They are hypotriploid and present an epithelial-like morphology. A375 cells form tumors when injected into immune-compromised mice, with high proliferation rate and invasiveness, and is useful for xenograft studies. It is widely used in mouse models of skin cancer. The presence of the luciferase reporter allows for easy assay readouts as luciferase activity is proportional to the number of cells.

Application

- Use as an internal control in co-culture killing assays
- *In vitro* and *in vivo* bioluminescence imaging

Materials Provided

Components	Format
2 vials of frozen cells	Each vial contains $>1 \times 10^6$ cells in 1 ml of Cell Freezing Medium (BPS Bioscience #79796)

Parental Cell Line

A375 cells, human melanoma cell line, epithelial cells, adherent.

Mycoplasma Testing

The cell line has been screened to confirm the absence of Mycoplasma species.

Materials Required but Not Supplied

These materials are not supplied with the cell line but are necessary for cell culture and cellular assays. BPS Bioscience's reagents are validated and optimized for use with this cell line and are highly recommended for best results. Media components are provided in the Media Formulations section below.

Media Required for Cell Culture

Name	Ordering Information
Thaw Medium 6	BPS Bioscience #60183
Growth Medium 6C	BPS Bioscience #78077

Materials Required for Cellular Assay

Name	Ordering Information
Thaw Medium 6	BPS Bioscience #60183
96-well tissue culture-treated white clear-bottom assay plate	
ONE-Step™ Luciferase Assay System	BPS Bioscience #60690
Luminometer	

Storage Conditions



Cells are shipped in dry ice and should immediately be thawed or stored in liquid nitrogen upon receipt. Do not use a -80°C freezer for long term storage. Contact technical support at support@bpsbioscience.com if the cells are not frozen in dry ice upon arrival.

Media Formulations

For best results, the use of validated and optimized media from BPS Bioscience is *highly recommended*. Other preparations or formulations of media may result in suboptimal performance.



Note: Thaw Media do *not* contain selective antibiotics. However, Growth Media *do* contain selective antibiotics, which are used to maintain selective pressure on the cell population expressing the gene of interest. Cells should be grown at 37 °C with 5% CO₂. BPS Bioscience's cell lines are stable for at least 10 passages when grown under proper conditions.

Media Required for Cell Culture

Thaw Medium 6 (#60183):

DMEM medium supplemented with 10% FBS and 1% Penicillin/Streptomycin.

Growth Medium 6C (#78077):

DMEM medium supplemented with 10% FBS, 1% Penicillin/Streptomycin, and 0.25 µg/ml Puromycin.

Media Required for Functional Cellular Assay

Thaw Medium 6 (#60183):

DMEM medium supplemented with 10% FBS and 1% Penicillin/Streptomycin.

Cell Culture Protocol

Note: A375 cells are derived from human material and thus the use of adequate safety precautions is recommended.

Cell Thawing

1. Swirl the vial of frozen cells for approximately 60 seconds in a 37°C water bath. As soon as the cells are thawed (it may be slightly faster or slower than 60 seconds), quickly transfer the entire contents of the vial to a tube containing 10 ml of pre-warmed Thaw Medium 6.

Note: Leaving the cells in the water bath at 37°C for too long will result in rapid loss of viability.

2. Immediately spin down the cells at 300 x g for 5 minutes, remove the medium and resuspend the cells in 5 ml of pre-warmed Thaw Medium 6.
3. Transfer the resuspended cells to a T25 flask or T75 flask and incubate at 37°C in a 5% CO₂ incubator.
4. After 24 hours of culture, check for cell attachment and viability. Change medium to fresh Thaw Medium 6 and continue growing in a 5% CO₂ incubator at 37°C until the cells are ready to passage.

- Cells should be passaged before they are fully confluent. At first passage and subsequent passages, use Growth Medium 6C.

Cell Passage

- Aspirate the medium, wash the cells with phosphate buffered saline (PBS) without $\text{Ca}^{2+}/\text{Mg}^{2+}$, and detach the cells from the culture vessel with 0.25% Trypsin/EDTA.
- Once the cells have detached, add Growth Medium 6C and transfer to a tube.
- Spin down cells at $300 \times g$ for 5 minutes, remove the medium and resuspend the cells in Growth Medium 6C. Seed into new culture vessels at the recommended sub-cultivation ratio of 1:2 to 1:8 once or twice per week.

Cell Freezing

- Aspirate the medium, wash the cells with PBS without $\text{Ca}^{2+}/\text{Mg}^{2+}$, and detach the cells from the culture vessel with 0.25% Trypsin/EDTA.
- Once the cells have detached, add Growth Medium 6C and count the cells.
- Spin down the cells at $300 \times g$ for 5 minutes, remove the medium and resuspend the cells in 4°C Cell Freezing Medium (#79796) at $\sim 2 \times 10^6$ cells/ml.
- Dispense 1 ml of cell suspension into each cryogenic vial. Place the vials in an insulated container for slow cooling and store at -80°C overnight.
- Transfer the vials to liquid nitrogen the next day for long term storage.



Note: It is recommended to expand the cells and freeze at least 10 vials at an early passage for future use.

Validation Data




Figure 1: Luciferase activity in the Firefly Luciferase A375 Cell Line.

Firefly Luciferase A375 cells were seeded into a 96-well plate at 5000 cells/well in 50 μl of Thaw Medium 6, and luciferase activity was measured using the ONE-Step™ Luciferase Assay System (#60690).

Data shown is representative. For lot-specific information, please contact BPS Bioscience, Inc. at support@bpsbioscience.com

License Disclosure

Visit bpsbioscience.com/license for the label license and other key information about this product.

Troubleshooting Guide

Visit bpsbioscience.com/cell-line-faq for detailed troubleshooting instructions. For all further questions, please email support@bpsbioscience.com.

Related Products

<i>Products</i>	<i>Catalog #</i>	<i>Size</i>
Firefly Luciferase Lentivirus	79692	500 µl x 2
eGFP/Firefly Luciferase U-87 MG Cell Line	78904	2 vials
eGFP/ Firefly Luciferase MM.1S Cell Line	78376	2 vials
eGFP/ Firefly Luciferase RS4;11 Cell Line	78926	2 vials
eGFP/Firefly Luciferase K562 Cell Line	78911	2 vials
eGFP/Firefly Luciferase OVCAR3 Cell Line	78953	2 vials

Version 121324