



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6405 Mira Mesa Blvd, Suite 100
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: support@bpsbioscience.com

Human Interferon-gamma Recombinant

Catalog #: 90162-B

Lot #: 230417

Description: Recombinant Human Interferon-gamma is a disulfide-linked monomeric protein consisting of 144 amino acids and migrates as an approximately 17 kDa protein under reducing and non-reducing conditions in SDS-PAGE. Optimized DNA sequence encoding Human Interferon-gamma mature chain was expressed in *E. coli*.

Source: Optimized DNA sequence encoding human Interferon-gamma mature chain was expressed in *E. coli*.

Formulation: Lyophilized from a 0.2 µm filtered concentrated (1mg/ml) solution in PBS, pH 7.2.

Reconstitution: A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/ml. This solution can then be diluted into other buffers.

Storage: The lyophilized protein is stable for at least 2 years from date of receipt when stored at -20°C. Upon reconstitution, store in working aliquots at 2 - 8° C for up to one month, or at -20°C for up to six months, in the presence of a carrier protein. Avoid repeated freeze/thaw cycles.

Purity: >98%, as determined by SDS-PAGE and HPLC.

Endotoxin Level: Endotoxin level was found to be < 0.1 ng/µg (1EU/µg), using the LAL gel clot method.

Biological Activity: The ED50, was determined by a viral resistance assay of mouse L929 cells and was found to be ≤ 1 ng/ml, corresponding to a specific activity of ≥ 1 x 10⁶ units/mg.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**

Or you can Email us at: support@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com