



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



6405 Mira Mesa Blvd., Ste 100
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: support@bpsbioscience.com

Human Interleukin-4 Recombinant

Catalog #: 90193-A
Lot#: 220722

Description: Recombinant human interleukin-4 (IL-4), also known as B-Cell Growth Factor (BCGF), is a disulfide-linked monomeric protein consisting of 130 amino acid residue subunits and migrates as an approximately 15 kDa protein under reducing and non-reducing conditions.

Source: Optimized DNA sequence encoding human IL-4 mature chain was expressed in *E. coli*.

Formulation: Lyophilized from a 0.2 µm filtered solution containing 2.5% glycine, 0.5% sucrose, 0.01% Tween-80, 5 mM glutamic acid, pH 4.5.

Reconstitution: A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/ml. This solution can then be diluted into other buffers.

Storage: The lyophilized protein is stable for at least 2 years from date of receipt when stored at -20°C. Upon reconstitution, store in working aliquots at 2 - 8°C for up to one month, or at -20°C for up to six months, in the presence of a carrier protein. Avoid repeated freeze/thaw cycles.

Purity: >95%, as determined by SDS-PAGE and HPLC.

Endotoxin Level: Endotoxin level was found to be < 0.1 ng/µg (1 EU/µg), using the LAL gel clot method.

Biological Activity: The ED₅₀, determined by the dose-dependent stimulation of the proliferation of human TF-1 cells, is ≤ 0.1 ng/ml, corresponding to a specific activity of > 1x 10⁷ units/mg.

Protein Sequence:

MGLTSQLLPPLFFLLACAGNFVHGHKCD
ITLQEIIKTLNSLTEQKTLCTELTVTDIFAA
SKNTTEKETFCRAATVLRQFYSHHEKDT
RCLGATAQQFHRHKQLIRFLKRLDRNL
WGLAGLNSCPVKEANQSTLENFLERL
KTIMREKYSK CSS

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.
To place your order, please contact us by Phone **1.858.202.1401** Fax **1.858.481.8694**
Or you can Email us at: support@bpsbioscience.com
Please visit our website at: www.bpsbioscience.com