

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



12-oxo Stearic Acid methyl ester

Item No. 10007012

CAS Registry No.: 2380-27-0

Formal Name: 12-oxo-octadecanoic acid, methyl ester

Synonyms: Methyl 12-oxo Octadecanoate,

Methyl 12-keto Stearate,


Methyl 12-oxo Stearate

MF: $C_{19}H_{36}O_3$ FW: 312.5 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

12-oxo Stearic acid methyl ester is supplied as a crystalline solid. A stock solution may be made by dissolving the 12-oxo stearic acid methyl ester in the solvent of choice, which should be purged with an inert gas. 12-oxo Stearic acid methyl ester is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 12-oxo stearic acid methyl ester in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 12-oxo stearic acid methyl ester can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 12-oxo stearic acid methyl ester in PBS, pH 7.2, is approximately 0.15 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

12-oxo Stearic acid methyl ester is an esterified form of 12-oxo stearic acid. It has been found in esterified L. chinensis and L. decipiens pulp oils and serves as a measure of 12-oxo stearic acid content. 1

Reference

1. Kadry, H., Shoala, S., El Gindi, O., et al. Chemical characterization of the lipophilic fraction of Livistona decipiens and Livistona Chinensis fruit Pulps (palmae) and assessment of their anti-hyperlipidemic and anti-ulcer activities. Nat. Prod. Commun. 4(2), 265-270 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 06/15/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM