

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

# SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## N-3-oxo-dodecanoyl-L-Homoserine lactone

Item No. 10007895

CAS Registry No.: 168982-69-2

Formal Name: 3-oxo-N-[(3S)-tetrahydro-2-oxo-3-furanyl]-

dodecanamide


Synonyms: N-(3-oxododecanoyl) L-Homoserine lactone,

3-oxo-C12-HSL

MF: C<sub>16</sub>H<sub>27</sub>NO<sub>4</sub> FW: 297.4 **Purity:** ≥98% UV/Vis.:  $\lambda_{max}$ : 250 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



#### **Laboratory Procedures**

N-3-oxo-dodecanoyl-L-Homoserine lactone (3-oxo-C12-HSL) is supplied as a crystalline solid. A stock solution may be made by dissolving the 3-oxo-C12-HSL in the solvent of choice, which should be purged with an inert gas. 3-oxo-C12-HSL is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 3-oxo-C12-HSL in these solvents is approximately 20 mg/ml.

3-oxo-C12-HSL is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 3-oxo-C12-HSL should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 3-oxo-C12-HSL has a solubility of approximately 0.2 mg/ml in a 1:3 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

3-oxo-C12-HSL is a bacterial quorum-sensing signaling molecule produced by P. aeruginosa and strains of the B. cepacia complex. 1,2 It induces the production of IL-8 in 16HBE human bronchial epithelial cells when used at a concentration of 100 µM.3 Supernatants from 3-oxo-C12-HSL-stimulated 16HBE cells induce chemotaxis of isolated human neutrophils. 3-oxo-C12-HSL has been found in respiratory secretions from patients with cystic fibrosis infected with P. aeruginosa.<sup>2</sup>

#### References

- 1. Pearson, J.P., Gray, K.M., Passador, L., et al. Structure of the autoinducer required for expression of Pseudomonas aeruginosa virulence genes. Proc. Natl. Acad. Sci. USA 91(1), 197-201 (1994).
- 2. Chambers, C.E., Visser, M.B., Schwab, U., et al. Identification of N-acylhomoserine lactones in mucopurulent respiratory secretions from cystic fibrosis patients. FEMS Microbiol. Lett. 244(2), 297-304
- 3. Smith, R.S., Fedyk, E.R., Springer, T.A., et al. IL-8 production in human lung fibroblasts and epithelial cells activated by the Pseudomonas autoinducer N-3-oxododecanoyl homoserine lactone is transcriptionally regulated by NF-kB and activator protein-2. J. Immunol. 167(1), 366-374 (2001).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/20/2021

### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM